

GEOLOGY
OF THE
WILLOW RIDGE ANTICLINE
WISLEY AREA, N.W.T.

PREPARED FOR
CANADACITIES SERVICE PETROLEUM CORP.

BY
W. E. BRADY CONSULTING LTD.

CANARY, ALBERTA

AUGUST, 1968

OCT 13 1966
RESOURCE MANAGEMENT DIVISION

G4C

GEOLOGY

of the

WILLOW RIDGE ANTICLINE

WRIGLEY AREA, N.W.T.

Abstracted for
Geo-Science Data Index

Date _____

Prepared for

Canada-Cities Service Petroleum Corp.

By

W. B. Brady Consulting Ltd.

Calgary, Alberta

August, 1966

APPROVED
OCT 12 1966
RESOURCE MANAGEMENT DIVISION

INTRODUCTION

This report consists of the presentation of the results of a one-week geological investigation in the Wrigley-Fort Simpson area. This operation was carried out on behalf of Canada-Cities Service Petroleum Corporation between May 22 and May 29, 1966 by W. B. Brady Consulting Ltd. of Calgary. A Bell Super-G helicopter, chartered from Bullock Wings and Rotors, was used throughout the operation.

The purpose of the investigation was to study the stratigraphy along the Willow Ridge anticline and to compare and correlate the exposures there with those exposed in the McConnell Range to the west. In addition, a Middle Devonian reef on the Root River was examined in order better to understand the nature of the isolated reefs which may be present in the Corporation's Interior Plains acreage.

Base camp was established at Wrigley airport where board and lodging were supplied by the airport manager, A. Silverton. Gasoline was purchased from the radio operator, J. Ito. The writer wishes to extend his thanks to these gentlemen and to the pilot, R. Crawford, and the geological assistant, D. Gardiner.

The area under discussion had already been mapped by the Geological Survey of Canada during Operation Mackenzie in 1957. No actual landings were made on the Willow Ridge anticline at that time.

INDEX MAP
NORTHWEST CANADA



However, the writer was most fortunate to have been a participant in Operation Icehouse and the regional experience acquired thus cannot well during this recent study.

A geological map at the scale of one mile to one inch is enclosed with this report. Fossil paleontological identifications of fossil collections are included in the Appendix.

Physical Features of the Area

The map-area lies within the western part of the Interior Plains and encompasses elements of the Frontian Mountains and the Subarctic Plains.

The elevations within the map-area range from less than 1,000 feet to more than 2,500 feet above sea level. Lakes of various sizes dot the Interior Plains and afford easy access for forest plants over much of the area. The Plains are heavily wooded and blanketed with glacial debris. Glacial lineaments are readily visible from air photographs and many of the lakes have their major shorelines parallel to a former direction of ice movement. Traces of two distinct ice movements are apparent, the more prominent, and probably the more recent, having a trend of $S\ 75^{\circ}\ W$, the less distinct trending $S\ 30^{\circ}\ W$. Variations occur in both lineations, however, due mainly to effects of relief on ice movements.

Aside from Willow Ridge itself, a narrow artificial barrier of resistant Middle Devonian rocks, outcrops are limited to the Willowlake River and the McManis Range.

The Franklin Mountains, of which the Russell Range forms the main element within the map-area, are composed of resistant carbonates which have been brought to the surface by west-dipping thrust faults. South of the area, in the vicinity of Mt. Cap, a thick sequence of Proterozoic dolines and sandstones are exposed.

The Permian Plains, which are occasionally broken by higher elevations such as the Russell Range and the San Flats, extend to the west of the Franklin Mountains, beyond the western border of the map-area.

Most of the surveys within the map-area has been extensively covered by seismic operations. Many seismic lines resulting from this work can be observed on air-photographs, although none of the work post-dates the photography.

The area is best suited for winter drilling or seismic work after the snowing is frozen. In preparation for the winter work, heavy equipment should be moved along the Redoubt River by barge during the summer months.

A telephone line is now in operation at Vigley, enabling one to speak directly to the "outside world". This line is being extended to Inuvik, with completion date being anticipated for the fall of 1966.

Scheduled air service to Vigley is only available twice monthly, although chartered planes are available from Hay River, Fort Reliance, or Inuvik.

STRATIGRAPHY

General

Approximately 2,000 to 3,000 feet of strata underlie the Plains to the east of the map-area. On the McConnell Range, some 6,000 feet of strata are present between the top of the Middle Devonian and the top of the Proterozoic. Further to the west, over 15,000 feet of Paleozoic strata are present. This estimate does not include the Upper Devonian and/or Cretaceous rocks, which are very similar in appearance and can be confused in the field.

The great change in thickness of section eastwards presents a potential situation for the occurrence of many natural stratigraphic traps for the containment of hydrocarbons.

The rocks examined in detail during this study were principally the Middle Devonian Nahanni and Arnica formations on the Willow Ridge anticline. Exposures of these formations were compared with equivalent strata on the McConnell Range, although in this latter area the Arnica equivalent is better referred to as Bear Rock formation because of the high degree of brecciation present. In addition to these beds, the Mount Kindle formation was examined along the southern part of the McConnell Range in order to be able to evaluate the eastward extension of this formation into the Plains more precisely.

Exposures of the shales of the Fort Simpson formation and of the Cretaceous are rare but these were examined wherever possible. This work was limited mainly by the availability of helicopter landing sites.

Other formations, which are part of the general regional stratigraphic sequence but which are not exposed in the immediate area, are discussed briefly in a general summary which is based upon published reports and the writer's previous knowledge of the area.

The following two tables, reproduced from Geological Survey of Canada Paper 62-33, provides a succinct outline of correlations in the Lower Paleozoic (Table 1) and in the Devonian (Table 2).

PALEOZOIC

Ordovician and Silurian

The Sombre, Cansell, Delorme, Whittaker and Sunblood formations are exposed to the west in the Root River map-area but are absent within the project area. The Franklin Mountain formation may be equivalent in part to the Sunblood formation, and the Mount Kindle formation of the McConnell Range may be equivalent to the Whittaker formation to the west.

The Mount Kindle was examined in the core of the anticline along the McConnell Range where it is cut by the River Between Two Mountains. Approximately 300 feet of strata assignable to the Mount Kindle was observed at this locality although the base of the formation was not exposed. The formation at this exposure consists of dark grey weathering, thick, irregularly bedded, very fine-grained dolomite. Irregular blebs and nodules of chert are rare to common, many of them representing silicified fossil fragments. In general the formation is quite tight with only a few isolated vugs. At other localities, however, the Mount Kindle is reefy and is quite porous.

	Camsell Bend and Wrigley Map-areas (Douglas and D.K. Norris, 1961)	Dahadinni River Map-area		Wrigley Map-area		Horn River Map-area (Douglas and A.W. Norris, 1960)
		Rouge, Redstone and Dusky Ranges	Dahadinni Range	Camsell Range	McConnell Range	
Overlying beds	Arnica (11)	Arnica (15)	Arnica (15)	Arnica (15)	Bear Rock (16)	(12)
Devonian(?)	Sombre (10)	Sombre (14)	Camsell(?) (13)	? (12) ?		
	Camsell (5)	Camsell (13)				
Silurian	Delorme (4)	Delorme (11)				
	Whittaker (3)	Whittaker (10)			— ? — Mt. Kindle (9)	— ? — (10)
Ordovician	Sunblood (1)				— ? — Franklin Mt. (8)	— ? — (9)
— ? —	— ? —	— ? —			— ? — Saline River (6)	— ? — (8)
Cambrian					Mt. Cap	(7)
					Mt. Clark (5)	?
Proterozoic		(7)			Lone Land (4)	
					(3)	
					(2)	
					(1)	

Figure 1. Correlation of Lower Palaeozoic formations of Dahadinni River, Wrigley and adjacent map-areas.
Table

The Mount Kindle formation probably correlates with G.S.C. map-unit 10 of the Horn River area and must underlie all of the acreage between the McConnell Range and the Precambrian Shield to the east. Taking into consideration the porosity observed at other localities, the Mount Kindle forms a potential reservoir over a large area.

Devonian

The Bear Rock formation is the oldest Devonian rock found in the project area. It is Middle Devonian in age with the possibility that the lower part ranges as low as Lower Devonian. It consists predominantly of massive breccias composed of angular fragments of limestone and dolomite enclosed in a calcareous matrix. The formation tends to weather into rounded knobs, a situation which has led some geologists into describing such outcrops as reefs. This type of weathering is well developed in the Franklin Mountains (Figure 13). The stratigraphic equivalent of this formation in the subsurface contains much anhydrite, which probably accounts for the brecciation observable in the surface exposures. The Bear Rock formation may be correlated regionally with the Chinchaga formation of northern Alberta and British Columbia.

In outcrop the Bear Rock gives the impression of being more porous than it actually is, due to the weathering out of many individual fragments to form angular vugs. Some interconnected porosity is undoubtedly present, however, as can be deduced from the wells drilled into the formation which have yielded either fresh or salt water on test.

The Bear Rock formation changes facies to the southwest into the more definitely bedded and banded dolomites of the Arnica formation, (formerly called the Lone Mountain formation in some of the older reports). This change can be seen to begin to take place in outcrops south of the River Between Two Mountains and is complete by the time Lone Mountain, an outlier at the northern end of the Cassell Range, is reached.

Willow Ridge lies within the transitional area of the change from the brecciated Bear Rock to the better bedded dolomites of the Arnica. As the strata exposed in the section measured on Willow Ridge are predominantly well bedded, the term 'Arnica' is the better to use for the rocks at this location.

The various abrupt facies changes within the Devonian which are present in the mountains to the west and to the southwest are not present in the map area. A most notable change takes place between the shales and highly argillaceous limestones of the Funeral formation and the coarse crystalline porous dolomites of the Manetoe formation. This change is very abrupt and is well displayed at two locations, one near the Ram River and the other near Iverson Lake. One of the most interesting and unexpected observations made on Willow Ridge is the presence of the "Manetoe facies" in the core of the anticline. The presence of this dolomite on Willow Ridge indicates a much more extensive distribution of this formation than had previously been suspected, enhancing the prospects for possible Manetoe production from additional acreage.

The Nahanni formation is considerably thinner over the map-area than are the equivalent strata to the west. Thickness variations may range from 956 feet in the Iverson Range to the west, to less than 300 feet in the McConnell Range and 242 feet on Willow Ridge. This variation in thickness is due partly to original depositional variations in thickness combined with possible facies changes at the top of the formation.

The Nahanni formation was examined on Willow Ridge where it consisted of micrograined, grey limestone with numerous brachiopods and stromatoporoids near the top. The lithology of this formation, where examined on the McConnell Range, appeared to be quite similar to this in character. The shale facies-equivalent of the Nahanni formation, the Headless formation, is not present in this area, being present only to the west.

A summary of the section exposed on Willow Ridge follows:

Section measured on Willow Ridge anticline at approximately 122° 53' W and 62° 47' N., (described in descending order).

Strata overlying the Nahanni formation not exposed.

Middle Devonian

Nahanni formation

Limestone, grey, micrograined, thin (2 - 4")
bedded, somewhat rubbly, fossil collection
5-66-A.

Unit Thickness

63.0

	<u>Unit Thickness</u>
Covered interval - apparently quite shaly.	25.0
Limestone, brownish-grey, micrograined, finely laminated, a few blebs of calcite - possibly algal.	68.0
Partly covered but appears similar to interval below.	35.0
Limestone, pale brown, grey, micrograined, thin to medium bedded	21.0
Covered interval.	30.0
Total thickness of Nahanni formation	242.0

Nahanni formation was measured approximately 3/4 mile south of the lower part of the section. The 12-foot dolomite bed was present at both localities.

Arnica formation

Dolomite, light grey, coarse-grained, coarsely laminated, porous, may vary in thickness along strike.	12.0
Dolomite, brownish-grey, finely granular, brownish-grey, weathering, irregularly bedded, locally brecciated near top.	163.0
Dolomite, light grey, coarse crystalline, vuggy in part.	55.0
Limestone, grey, micrograined, grey weathering.	5.0
Dolomite, white to mottled, coarse crystalline, massive, faint traces of stromatoporoids.	10.0
Dolomite, grey to brownish-grey, finely granular, brownish-grey weathering, bedding planes irregular, calcite veins common, locally brecciated.	200.0

Total thickness of exposed
Arnica formation

445 feet

Total thickness measured

687 feet

The Nahanni formation is overlain by the Fort Simpson formation which consists of shale which is non-calcareous, dark grey, fissile and locally iron-stained. The upper part of this formation consists of medium to dark grey mudstone which is calcareous and silty and is interbedded with fine-grained argillaceous limestone and thinly bedded, laminated, fine-grained calcareous siltstone and sandstone. The lower non-calcareous unit of the formation may correlate with the Horn River formation.

The Fort Simpson formation was examined on the Willowlake River where it has been compressed into gentle undulating folds.

Numerous other Upper Devonian units which occur west of the map-area are not present to the east. One of these (G.S.C. map-unit 20 in the Camell Bend-Root River area, or G.S.C. map-unit 25 in the Dahadimi-Wrigley area) is of some interest in that it is composed largely of stromatoporoids and reefy coralline limestone beds which occur as isolated patches enclosed by the surrounding shale.

Cretaceous

Much of the Interior Plains is underlain by strata of Cretaceous age. They are shown on the geological map as flanking the Willow Ridge anticline, being separated from the Middle Devonian carbonates by the Fort Simpson shales. In the project area however, there appears to be no

distinctive mappable horizon which can be used to separate the Cretaceous from the Fort Simpson formation. Further to the south, the contact is placed at the base of a resistant ridge, although there has not been sufficient sampling of the outcrop to be certain that this is the actual boundary. It should be recognized therefore that although much of the Interior Plains to the east is indicated as being underlain by Cretaceous, at least part of the area presently mapped as Cretaceous may prove to be of Upper Devonian age. Detailed field work combined with micropaleontological determinations would be required for the solution of this problem.

POTENTIAL RESERVOIRS

The porous strata of the Sombre and Delorme formations which are exposed in the mountains west of the map-area are not present on the McConnell Range or on Willow Ridge. These formations may lie in a favourable structural position for the accumulation of hydrocarbons to the west of the map-area but should not be considered as potential horizons to the east of the area.

The Mount Kindle formation is a possible reservoir over much of the Interior Plains and is also present in the Horn River area (assuming that the correlation with G.S.C. map-unit 10 in that area is correct).

It should be noted at this point that it would be more accurate to relate all Middle Devonian terminology to facies as, in many cases, two distinct facies are found to be interbedded and it then becomes difficult to apply standard formation names. This situation occurs in the Nahanni

Range where Douglas has mapped the Landry formation, a micrograined limestone equivalent of the Manetoe dolomites. In actual fact both the Landry and the Manetoe facies are present jointly over most of the Nahanni Range.

The presence of the "Manetoe-type" facies is extremely interesting. These coarse-grained porous dolomites are obviously more widespread in development than had previously been suspected, increasing the potential area over which a reservoir may be anticipated.

What may become the main reservoirs in the area, however, are the isolated Devonian reefs similar to that present on the Horn River. At this latter locality the reef consists of massive, very coarse-grained, light to medium brown, richly fossiliferous limestone exposed on a circular hill 1/4 mile in diameter. These reefoid beds grade upwards into massive, pale brown, fine-grained limestone. A similar reef occurs on the Root River at Longitude $125^{\circ} 15'$. At this locality the reef forms a massive exposure (Figure 14) which consists of coarse-grained, crinoidal limestone in the lower part and fine-grained, stromatoporoidal limestone in the upper part. The reef is over 300 feet thick and extends for nearly two miles along strike. Its total lateral extent down-dip is presently unknown. The reef passes horizontally into shales which are eroded to form a broad valley. Brecciated, red and green shales are present between the reef and the Nahanni formation, which outcrops near the northern end of the reef. These shales contain much anhydrite together with what appears to be specular hematite, a possible indication of back-reef evaporitic deposits.

Fossils collected at the top of the Nahanni formation below the base of the reef on the Root River include:

Proetus spp.
Spinatrypa sp.
Ambocoelia sp.
Thamnopora sp.
Atrypa spp. aff. arctica Warren
Warrenella sp.
'Hypothyridina' aff. cameroni Warren
Productella sp.
Schizophoria sp.
Small rugose coral
Crinoid ossicles - abundant
Sponge spicules
Gastropod fragments
Small pelecypod.

Near the base of the reef, the following fossils were collected:

cf. Stringocephalus axius Crickmay
Large rugose coral fragment
Stromatoporoid - indet
cf. Atrypa sp.
Crinoid ossicles.

Near the top of the reef, a collection yielded the following:

Stringocephalus burtini Defrance in de Blainville
Atrypa sp.

Spinatrypa sp.

Rugose coral fragment

Crinoid ossicles - abundant

Stromatopora fragments.

This upper collection indicates a correlation with the Kee Scarp reef of the Hornum Wells area. The other collections, although indicating a Middle Devonian age, are not sufficiently diagnostic to indicate if an unconformity exists between the reef and the Nahanni formation, nor if the reef actually represents continuous sedimentation from the top of the Nahanni through the Hare Indian to the Kee Scarp equivalent.

As the reefs are isolated features, with only the two known locations at Horn and Root Rivers respectively, no trend along which other reefs may occur can be predicted with any reliability. The mere presence of the reef on Root River, however, indicates that other similar structures may occur under the Corporation's acreage on the Interior Plains.

An interesting feature was observed in an area south of the Willowlake River at approximately $121^{\circ} 35' W$ and $63^{\circ} 20' N$, following a study of air photographs. This feature consisted of a series of concentric (oval) and radial fractures forming a pattern approximately six miles long and three miles wide, trending east-west. Such a pattern might well be the surface reflection of one of the isolated reefs mentioned above. In any event, it is strongly recommended that this, and any similar features which may be located, be investigated in any proposed seismic work in the area.




PHOTO ANOMOLY

WILLOWLAKE RIVER

N.W.T.

Scale: Approximately 1"= 0.7 miles

RADIAL AND CONCRETIC LINES FROM AIR
PHOTOGRAPH PROBABLY REPRESENT FRACTURE
PATTERN.

W.B. Brady, Consulting Ltd.

STRUCTURE

Folding has been the most prominent result of tectonic activity within the map-area. The two main anticlines observed are the Willow Ridge anticline and the anticline along the McConnell Range.

The Willow Ridge anticline, (Figures 3, 4 and 10) approximately 28 miles long, forms a doubly-plunging fold which brings resistant Middle Devonian rocks to the surface. Dips off the flanks of the structure range from 30 to 75 degrees. The widest band of Middle Devonian rocks occurs in the southern part of the fold, where the outcrop is two miles wide. The main part of the anticline is narrow and rarely exceeds a width of more than one-half mile. The northerly plunge of the structure is gentle and the fold dies out in a series of small en-echelon anticlines. The main anticline plunges more steeply to the south, however, and has no obvious expression in the shales immediately to the south on Willowlake River. These shales are folded into a series of smaller folds, a phenomenon which is quite common where less competent beds overlie tightly folded competent beds.

The part of McConnell Range which lies within this map-area is also folded into an anticline (Figure 12) which is strongly asymmetric to the east. Dips on the west flank of the structure range from 25 to 35 degrees compared with east flank dips which range up to the vertical. Such a situation suggests the presence of a west dipping thrust fault on the east side of the structure.

Other folds similar to the Willow Ridge anticline may occur to the east beneath the western part of the Interior Plains. Such folds, if present, should be easily definable by seismic investigation.

CONCLUSIONS

As already mentioned the main objective of this field work was to establish the age and lithology of the strata exposed in the core of the Willow Ridge anticline and to effect a correlation between these beds and adjacent sections. A section measured in the most critical position for obtaining the maximum amount of information indicated that the strata consisted of elements of the Arnica and Nahanni formations. Out of a total of 687 feet of section measured, 445 feet were assigned to the Arnica and 242 feet to the Nahanni. The most interesting new observation, however, was the detection of the presence of the Manetoe facies within the Arnica bedded dolomites at this locality, indicating a wider distribution of this facies than had previously been thought to exist. Time permitted the investigation of an unmapped Middle Devonian reef on the Root River. The presence of this reef, which may correlate with that exposed on the Horn River, indicates the possibility of the existence of similar reefs (with sufficient cover to be of economic interest) beneath the Corporation's acreage on the Interior Plains.

A photo-anomaly consisting of concentric and radial fracture patterns may represent the surface expression of a buried reef such as mentioned above. It is recommended that any proposed seismic program should be planned so as to investigate such photo-anomalies thoroughly.

Finally, it is recommended that a regional geological study be undertaken which combines all subsurface data with published and unpublished surface information, a study which does not appear to have been carried out in this region by anyone to date. With regard to unpublished data, a little-known fact is that the Geological Survey of Canada in Ottawa maintains, on open file, unpublished descriptions of sections measured during Operation Mackenzie. Samples to accompany these section descriptions are also available for examination. Such material would form an excellent basis for the regional study mentioned above.

Respectfully submitted,

W. B. Brady
W. B. Brady, P. Geol.



Figure 3

View looking northwest through the gap near the southern part of the Willow Ridge anticline. The McConnell Range can be seen through the gap to the west.



Figure 3

View looking northwest through the gap near the southern part of the Willow Ridge anticline. The McConnell Range can be seen through the gap to the west.



Figure 4 View looking north of the gap shown in Figure 3. Top of lower cliff is capped by 12-foot bed of coarse-crystalline dolomite at the top of the Arnica formation. The upper cliff and covered interval between is the Nahanni formation.



Figure 4 View looking north of the gap shown in Figure 3. Top of lower cliff is capped by 12-foot bed of coarse-crystalline dolomite at the top of the Arnica formation. The upper cliff and covered interval between is the Nahanni formation.



Figure 5

Section measured in core of Willow Ridge anticline. There is more section exposed here than at the gap as shown in Figure 4. The lower coarse crystalline dolomite is outlined followed by very fine-grained dolomite and finally overlain by the upper 12-foot unit of coarse dolomite. The Nahanni formation was measured approximately one-mile south.




Figure 5

Section measured in core of Willow Ridge anticline. There is more section exposed here than at the gap as shown in Figure 4. The lower coarse crystalline dolomite is outlined followed by very fine-grained dolomite and finally overlain by the upper 12-foot unit of coarse dolomite. The Nahanni formation was measured approximately one-mile south.



Figure 6

View looking north along the Willow Ridge anticline immediately north of section shown in Figure 5. The upper light grey coarse-crystalline dolomite is overlain on the far right by the Nahanni formation.




Figure 6

View looking north along the Willow Ridge anticline immediately north of section shown in Figure 5. The upper light grey coarse-crystalline dolomite is overlain on the far right by the Nahanni formation.



Figure 7

Upper coarse crystalline dolomite (here thicker than where section was measured) in right foreground. The Nahanni formation can be seen through the trees in left foreground. The McConnell Range and the gap formed by The River Between Two Mountains can be seen in the background.



Figure 7

Upper coarse crystalline dolomite (here thicker than where section was measured) in right foreground. The Nahanni formation can be seen through the trees in left foreground. The McConnell Range and the gap formed by The River Between Two Mountains can be seen in the background.



Figure 8

Close-up view of the coarse-crystalline Manetoe "type-facies". Note porosity parallel to the rudimentary bedding, probably representing solution of fossil fragments.



Figure 8

Close-up view of the coarse-crystalline Manetoe "type-facies". Note porosity parallel to the rudimentary bedding, probably representing solution of fossil fragments.



Figure 8

Close-up view of the coarse-crystalline Manetoe "type-facies". Note porosity parallel to the rudimentary bedding, probably representing solution of fossil fragments.



Figure 9 Irregular contact between the coarse-crystalline dolomite on the right and Nahanni-type limestone on the left.




Figure 9 Irregular contact between the
coarse-crystalline dolomite on
the right and Nahanni-type lime-
stone on the left.



Figure 10

View looking north along Willow Ridge anticline. This structure, especially in this northern part is very tight and plunges northward in a series of en echelon folds. It is mainly tree covered affording few helicopter landing locations.




Figure 11

Low dipping beds of the Nahanni formation near the south end of Old Fort Island on the Mackenzie River. View is looking northwest with the Camell Range in the distance.




Figure 11

Low dipping beds of the Nahanni formation near the south end of Old Fort Island on the Mackenzie River. View is looking northwest with the Camsell Range in the distance.




Figure 12

View looking north along the McConnell Range with the main part of the Franklin Mountains in the distance. This picture shows clearly the anticlinal nature of the McConnell Range. The gap formed by the River Between Two Mountains is immediately to the north. The upper coarse-crystalline dolomite, as on the Willow Ridge anticline is in the left foreground underlain here by Bear Rock. The Mount Kindle formation can be seen in the core of the anticline with the Nahanni formation on the flank to the left.



Figure 12

View looking north along the McConnell Range with the main part of the Franklin Mountains in the distance. This picture shows clearly the anticlinal nature of the McConnell Range. The gap formed by the River Between Two Mountains is immediately to the north. The upper coarse-crystalline dolomite, as on the Willow Ridge anticline is in the left foreground underlain here by Bear Rock. The Mount Kindle formation can be seen in the core of the anticline with the Nahanni formation on the flank to the left.



Figure 13

Knobby weathering Bear Rock formation immediately north of Figure 14. On the Willow Ridge anticline the same stratigraphic interval would be called Arnica as it is better bedded.



Figure 13

Knobby weathering Bear Rock formation immediately north of Figure 14. On the Willow Ridge anticline the same stratigraphic interval would be called Arnica as it is better bedded.



Figure 14

Massive biohermal reef overlying the Nahanni formation on Root River. The lower part of this reef is coarse-grained crinoidal limestone and the upper part is composed of *Stromatipora* limestone with "Stringocephalus".



Figure 14

Massive biohermal reef overlying the Nahanni formation on Root River. The lower part of this reef is coarse-grained crinoidal limestone and the upper part is composed of *Stromatipora* limestone with "Stringocephalus".

Collectors Initials or Name 1-66-C	Loc. Number	Dist. and Year	Datum and Footage from Datum		Catalog Number	MACRO MICRO	
Outcrop or Well Name		Location			Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
Biota and remarks: Large stromatoporeid 'heads' including: Clathradictyon sp. cf. Stromatopora sp.							
Stage	Biozone or Strat. equivalents		Ident. by	Date	Card of		
	Cambrian - Permian		D.W.	June/66	Lithology of Interval		
Validity of age determination (Checked)				ENVIRONMENT			
☒ Good ☐ Fair ☐ Poor							

Collectors Initials or Name 3-66-A	Loc. Number	Dist. and Year	Datum and Footage from Datum		Catalog Number	MACRO MICRO	
Outcrop or Well Name		Location			Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
Biota and remarks: cf. Devonogypa sp. (Gypidula sp. A. & McLaren) Hexanema sp. Small gastropod cross-sections Crinoid ossicles Ostracoda cross-sections Camarites sp.							
Stage	Biozone or Strat. equivalents		Ident. by	Date	Card of		
	M Devonian		D.W.	June/66	Lithology of Interval		
Validity of age determination (Checked)				ENVIRONMENT			
☒ Good ☐ Fair ☐ Poor							

Collectors Initials or Name S-16-A		Loc. Number	Dist. and Year	Datum and Footage from Datum		Catalog Number	MACRO MICRO
Outcrop or Well Name			Location		Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
Biotite and remarks: <i>Schuchertella adacta</i> Crickmay <i>Dorvillina</i> sp. <i>Productella</i> sp. - fragments <i>Hexagonaria</i> sp. <i>Spirifer</i> sp. - indet. Graptolite colonies including typical <i>Hume</i> forms. Productid fragments Graptolite fragments							
Stage		Biozone or Strat. equivalents		Ident. by	Date	Lithology of Interval	
Carbonian		Hume		DW.	June/66		
Validity of age determination (Checked) ☒ Good ☐ Fair ☐ Poor				ENVIRONMENT 			


Collectors Initials or Name 6-66-A		Loc. Number	Dist. and Year	Datum and Footage from Datum		Catalog Number	MACRO MICRO
Outcrop or Well Name			Location		Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
Biotite and remarks: <i>Hexagonaria</i> sp.							
Stage		Biozone or Strat. equivalents		Ident. by	Date	Lithology of Interval	
		Devonian		DW	June/66		
Validity of age determination (Checked) ☒ Good ☐ Fair ☐ Poor				ENVIRONMENT 			


Collector's Initials or Name 8-66-A		Loc. Number	Dist. and Year	Datum and Footage from Datum	Catalog Number	MACRO MICRO	
Outcrop or Well Name			Location		Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
Biotite and remarks: Proetus spp. - pygidia and fragments of cephalon. Spinatrypa sp. - fragments Ambocoelia sp. Thamnospira sp. Small rugose coral - fragments Small pelymptid - indet. Crinoid (fragments) - some aff. Leptomena sp. Small spicules. Crinoid ossicles							
Stage	Biozone or Strat. equivalents			Ident. by	Date	Card	of
	Devonian - probably Middle Devonian			D.W.	June/66		
Validity of age determination (Checked)				ENVIRONMENT			
☒ Good ☐ Fair ☐ Poor							


Collector's Initials or Name 8-66-B.		Loc. Number	Dist. and Year	Datum and Footage from Datum	Catalog Number	MACRO MICRO	
Outcrop or Well Name			Location		Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
Biotite and remarks: Atrypa spp. aff. arctica Warren. Spinatrypa spp. Warrenella sp. Ambocoelia sp. Thamnospira sp. Hypothyridina aff. cameroni Warren. Productella sp. - fragments Schizophoria sp. - fragments. Small rugose coral Crinoid ossicles - abundant.							
Stage	Biozone or Strat. equivalents			Ident. by	Date	Card	of
	M Devonian.			D.W.	June/66		
Validity of age determination (Checked)				ENVIRONMENT			
☒ Good ☐ Fair ☐ Poor							

Collectors Initials or Name 8-66-C		Loc. Number	Dist. and Year	Datum and Footage from Datum		Catalog Number	MACRO MICRO
Outcrop or Well Name			Location		Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
State and remarks: cf. <i>Stringocephalus</i> <i>adina</i> <i>Crickmay</i> <i>large ridged corals - fragment</i> <i>Stromatopora</i> - <i>indet.</i> cf. <i>Atrypa</i> sp. <i>Crinoid ossicles.</i>							
Card 1 of							
Stage	Biozone or Strat. equivalents		Ident. by	Date	Lithology of Interval		
	M. Devonian.		D.W.	June/66			
Validity of age determination (Checked)				ENVIRONMENT			
☐ Good ☒ Fair ☐ Poor							

Collectors Initials or Name 15		Loc. Number	Dist. and Year	Datum and Footage from Datum		Catalog Number	MACRO MICRO
Outcrop or Well Name			Location		Formation Name		
(Field) Geologist	Date	Notebook	Page	Field Paleontologist	Date	Notebook	Page
State and remarks: <i>Stringocephalus burtini</i> <i>Defrance in de Blinville.</i> <i>Atrypa</i> sp. <i>Spinatrypa</i> sp. <i>large corals - fragment.</i> <i>Crinoid ossicles - abundant.</i> <i>Stromatopora</i> <i>fragments.</i>							
Card of							
Stage	Biozone or Strat. equivalents		Ident. by	Date	Lithology of Interval		
Givetian	Kee Scarp.		D.W.	June/66			
Validity of age determination (Checked)				ENVIRONMENT			
☒ Good ☐ Fair ☐ Poor							







Base Map Control from Canada, Mills Lake Quad (E/2) 1:250,000 Sheet 85E
Photography - RCAF 1947, 1948, 1950 & 1951

Cities Service Aroage

Local Photo Anomaly
Regional Photo Anomaly

EXPLANATION


QUATERNARY	Qal	Aluvium
	Qls	Beach Deposits and Lake Sills
	Qgm	Glacial Moraine Deposits
CRETACEOUS	Kfn	Fort Nelson
	Kfsj	Fort St. John
DEVONIAN	Dtr	Trout River
	Dk	Kakisa Ls.
	Dgm	Grumbler, Upper
	Dgm	Grumbler, Middle
	Da	Alexandra Fm.
	Dh	Hay River Sh. & Ls.
	Dsp	Slope Point (?) Reef
	Dhr	Horn River Fm., Sh. & Ls. (Presqu'ile)
	Dpp	Pine Point Fm.




CITIES SERVICE OIL CO.



PRELIMINARY PHOTOGEOLOGIC MAP CANADA, NORTHWEST TERRITORIES Ft. Providence Project Mills Lake Quad E/2

Dip Component
 Dip Strike
 Topographic Slope
 Faults
 Shaded Relief
 Elevation
 Vegetation
 Bluffs
 Troughs
 Chert
 Surface Color - Li-Gy (Light Grey)









WILLOW RIDGE ANTICLINE


WRIGLEY — CAMSELL BEND AREA
N.W.T.

PREPARED FOR
CANADA — CITIES SERVICE PETROLEUM CORP.

BY
W.B. Brady, Consulting Ltd.

Calgary, Alberta.

AUGUST, 1966




MAP UNITS

- PLEISTOCENE AND PRESENT**
- 7 Alluvial sands and silts of the Mackenzie River.
- CRETACEOUS**
- 6 Shale, dark grey.
- UPPER (AND MIDDLE ?) DEVONIAN**
- 5 FORT SIMPSON FORMATION:
Shale and mudstone, grey.
- MIDDLE DEVONIAN**
- 4 NAHANNI FORMATION:
Limestone, micrograined, grey.
 - 3 ARNICA FORMATION:
Dolomite, grey bedded and banded.
 - 2 BEAR ROCK FORMATION:
Breccia of limestone and dolomite; facies equivalent of the Arnica fm.
- PRE-DEVONIAN**
- 1 MOUNT KINDE FORMATION:
Dolomite, grey, massive.

LEGEND

- Field Observed
- Helicopter Observed
- Published
- Vertical
- Overtuned
- Dip component
- Photogeologic dip, 45° to nearly vertical
- Photogeologic dip, 25° to 45°
- Photogeologic dip, 10° to 25°
- Photogeologic dip, 3° to 10°
- Geologic boundary (Defined, Approximate, Assumed)
- Fault



MAP UNITS

- PLEISTOCENE AND PRESENT**
- 7 Alluvial sands and silts of the Mackenzie River.
- CRETACEOUS**
- 6 Shale, dark grey.
- UPPER (AND MIDDLE ?) DEVONIAN**
- 5 FORT SIMPSON FORMATION:
Shale and mudstone, grey.
- MIDDLE DEVONIAN**
- 4 NAHANNI FORMATION:
Limestone, micrograined, grey.
- 3 ARNICA FORMATION:
Dolomite, grey bedded and banded.
- 2 BEAR ROCK FORMATION:
Breccia of Limestone and dolomite; facies equivalent of the Arnica fm.
- PRE-DEVONIAN**
- 1 MOUNT KINDLE FORMATION:
Dolomite, grey, massive.

LEGEND

- Field Observed.
- Helicopter Observed.
- Published.
- Vertical.
- Overturned.
- Dip component.
- Photogeologic dip, 45° to nearly vertical.
- Photogeologic dip, 25° to 45°.
- Photogeologic dip, 10° to 25°.
- Photogeologic dip, 3° to 10°.
- Geologic boundary (Defined, Approximate, Assumed).
- Fault.
- Thrust fault (Teeth in direction of dip).
- Anticline, - Semi-circle denotes plunge.
- Syncline, - Semi circle denotes plunge.
- Outcrop area.
- Ground observation with station number.
- Trend of glacial lineation.
- Measured section.
- Distinct Alignment

CANADA
MINES AND TECHNICAL SURVEYS
SURVEYS AND MAPPING BRANCH




PHOTO INDEX
PHOTO PROJECT
PHOTOS WORKED
MAP COMPLETE


Project
① Ft. Simpson
② Ft. Liard
③ Ft. Providence

INDEX TO 1:50,000 AND 1:63,360 MAP SHEETS
OF THE NATIONAL TOPOGRAPHIC SYSTEM

CORRECTED TO MAY 31, 1959

When ordering maps, the map name and number should be quoted.
EXAMPLE: - HAY RIVER 85 85 1/2
Map name 85 85 1/2
Block number 85
Letter B

MAPS PUBLISHED INDICATED THUS
1:50,000
1:50,000 Preliminary Edition
(incomplete as to colour, place names, etc.)
1:50,000 Advance Print
(print of manuscript one colour only)
P indicates an unprinted map



Published maps may be obtained from the Map Distribution Office, Department of Mines and Technical Surveys, Ottawa, Ont., at 25 cents per sheet. Maps at 1:50,000 scale are published in two sheets (E3 & W3) south of latitude 62°. Remittance should accompany order.

8-30-66

CANADA
MINES AND TECHNICAL SURVEYS
SURVEYS AND MAPPING BRANCH




PHOTO INDEX
PHOTO PROJECT
PHOTOS WORKED
MAP COMPLETE

Project
① Ft. Simpson
② Ft. Liard
③ Ft. Providence


INDEX TO 1:50,000 AND 1:63,360 MAP SHEETS
OF THE NATIONAL TOPOGRAPHIC SYSTEM

CORRECTED TO MAY 31, 1959

When ordering maps, the map name and number should be quoted.
EXAMPLE: - HAY RIVER 85 8/1


Map name
Sheet number
Letter



MAPS PUBLISHED INDICATED THUS:
1:50,000
1:50,000 Preliminary Edition
(incomplete as to colour, place names, etc.)
1:50,000 Advance Print
(print of manuscript or one colour only)
P indicates an uncoloured map




Published maps may be obtained from the Map Distribution Office, Department of Mines and Technical Surveys, Ottawa, Ont., at 25 cents per sheet. Maps at 1:50,000 scale are published in two sheets (E & W) 1. south of Latitude 62°.


8-30-66








11a2




Local Photo Assembly

[illegible]



Topographic Sheet (1:50,000) Sheet 85C
Photoreduced - 1:250,000 1960, 1965, 1970



Map of the District of Mackenzie, Alberta, showing the area around the town of Fort McMurray. The map is oriented horizontally and shows a complex network of ridges and valleys. Several large, dark, irregular shapes are overlaid on the map, possibly representing specific areas of interest or damage.


Legend


Scale

North Arrow

Index

1242





Well Abbreviations

SAHS JMC	British American - Hudson's Bay	Jean Marie Creek
STsLI	Briggs	Turkey Lake
W3A	Weston	3A
W7A	Weston	7A

Local Photo Anomaly ☒

Regional Photo Anomaly ☐

EXPLANATION



Div
Total Base Formula
Div


Gravelly
Formation


CITY'S SERVICE OIL CO.

PRELIMINARY PHOTOGEOLOGIC MAP
CANADA, NORTHWEST TERRITORIES
Ft. Simpson Quadrangle

[illegible][illegible]

English: 100%





Well Abbreviations

BABE JMC	British American - Hudson's Bay - Jean Marie Co
STULI	Stigge - Turkey Lake
WSA	Western SA
WTA	Western TA

Local Maps Assembly 

Regional Maps Assembly 

EXPLANATION

Quartzite facies

Q
Quartzite facies

Qtz
Late Quartzite

Gnl
Albite

Gg
Grossular

Gpm
Grossular


Gms
Grossular


STIMULI

[On]	Treat Stage Formation
[Off]	Initial Lb
[On]	Growth Formation
[Off]	Steady State

[illegible]







Well Abbreviations

BAHB JMC	British American - Hudson's Bay Jean Marie Creek
BTuLi	Briggs Turkey Lake
W3A	Westerol 3A
W7A	Westerol 7A

Local Photo Anomaly
Regional Photo Anomaly

EXPLANATION

Q	Undifferentiated
Qls	Lake Deposits
Qol	Alluvium
Qg	Glacial Undifferentiated
Qgm	Glacial Moraine

Kfn	Fort Nelson Formation
Kfsj	Fort St. John Group

Dtr	Trout River Formation
Dk	Kakisa Lt.
Dg	Grumbler Formation
Ds	Simpson Group




CITIES SERVICE OIL CO.

PRELIMINARY PHOTOGEOLOGIC MAP CANADA, NORTHWEST TERRITORIES Ft. Simpson Quadrangle NO. 1

Map Commission	DATE	SCALE
Map Title	1955	1:50,000
Map Scale	1:50,000	1:50,000
Map Scale	1:50,000	1:50,000
Map Scale	1:50,000	1:50,000
Map Scale	1:50,000	1:50,000

Graphic Scale





EXPLANATION

The diagram illustrates the geological column with three main periods: Quaternary, Cretaceous, and Devonian. Each period is associated with specific formations or deposits, some of which are grouped into sub-periods like Upper and Middle Devonian.

Period	Formation/Deposit	Location/Notes
QUATERNARY	Qol	
	Alluvium	
	Qls	
	Qgm	Beach Deposits and Lake Dike
CRETACEOUS	Kfn	Fort Nelson
	Kfs	Fort St. John
DEVONIAN	Dtr	Trout River
	Dk	
	Kakas Ls.	
	Dgo	
	Grumbler, Upper	
	Dgm	
	Grumbler, Middle	
	Ds	
	Alexandra Fm.	
	Dh	
	Hay River	Sh. & Ls.
	Dap	
	Steele Point (?) Reef	
	Dlr	
Horn River Fm., Sh. & Ls. (Fraserside)		
Dgo		
Flow Water Sh.		

Additional groupings and locations shown in the diagram:


- Quaternary Grouping:** Qol, Alluvium, Qls, Qgm (Beach Deposits and Lake Dike).
- Cretaceous Grouping:** Kfn (Fort Nelson), Kfs (Fort St. John).
- Devonian Grouping:**
 - Upper Devonian:** Dtr, Dk, Kakas Ls., Dgo, Grumbler, Upper, Dgm, Grumbler, Middle, Ds, Alexandra Fm., Dh.
 - Middle Devonian:** Hay River, Dap, Steele Point (?) Reef, Dlr, Horn River Fm., Sh. & Ls. (Fraserside), Dgo, Flow Water Sh.
- Other Locations:** Dfs (Fort Simpson) is associated with the Middle Devonian group.


CITIES SERVICE OIL CO

PRELIMINARY PHOTOGEOLOGIC MAP
CANADA, NORTHWEST TERRITORIES
Ft. Providence Project
Rise Quad SW/4

[illegible]

Copyright © 2004 John Wiley & Sons, Ltd.





Base Map Control from Canada, Ft. Liard Quad. 1:250,000 Sheet 958
Photography - Bradley 1961 1:60,000
Hunting 1964 1:60,000

WELL DATA LEGEND
Pb - Permian Belly
M - Mississippian Debol
M - Mississippian Bonff
Dw - Devonian Wabamun
Dsl - Devonian Simpson
Dap - Devonian Slave Point
Dp - Devonian Elk Point
Dc - Devonian Chicago
pre-Dev - pre-Devonian

Local Photo Anomaly

QUATERNARY

Qal Stream Alluvium
Qls
Landslide
Qo
Outwash deposits
Qg
Glacial Undifferentiated

CRETACEOUS

Kfn Fort Nelson Fm.
Kfsj Fort St. John Gp.
Kfs
Kfsi
Kfsk
Kfsl
Kfsm
Kfsn
Kfsr
Kfsu
Kfsv
Kfsx
Kfsy
Kfsz
Kfs1
Kfs2
Kfs3
Kfs4
Kfs5
Kfs6
Kfs7
Kfs8
Kfs9
Kfs10
Kfs11
Kfs12
Kfs13
Kfs14
Kfs15
Kfs16
Kfs17
Kfs18
Kfs19
Kfs20
Kfs21
Kfs22
Kfs23
Kfs24
Kfs25
Kfs26
Kfs27
Kfs28
Kfs29
Kfs30
Kfs31
Kfs32
Kfs33
Kfs34
Kfs35
Kfs36
Kfs37
Kfs38
Kfs39
Kfs40
Kfs41
Kfs42
Kfs43
Kfs44
Kfs45
Kfs46
Kfs47
Kfs48
Kfs49
Kfs50
Kfs51
Kfs52
Kfs53
Kfs54
Kfs55
Kfs56
Kfs57
Kfs58
Kfs59
Kfs60
Kfs61
Kfs62
Kfs63
Kfs64
Kfs65
Kfs66
Kfs67
Kfs68
Kfs69
Kfs70
Kfs71
Kfs72
Kfs73
Kfs74
Kfs75
Kfs76
Kfs77
Kfs78
Kfs79
Kfs80
Kfs81
Kfs82
Kfs83
Kfs84
Kfs85
Kfs86
Kfs87
Kfs88
Kfs89
Kfs90
Kfs91
Kfs92
Kfs93
Kfs94
Kfs95
Kfs96
Kfs97
Kfs98
Kfs99
Kfs100

PERMIAN & CARBONIFEROUS

P9
chert and ss.
CGS Bull. 59-6
Pm
Mottson Fm.


MISSISSIPPIAN


M7
Gym, Gy sh. and Gy ss.
CGS Bull. 59-6



CITIES SERVICE OIL CO. PRELIMINARY PHOTOLOGIC MAP CANADA, NORTHWEST TERRITORIES FT. LIARD QUADRANGLE NO. 1

Map Component
Date
Geology
Plotting
Drafting
Checked
Scale
Graphic Scale







City Service Acreage
Local Photo Anomaly

EXPLANATION

QUATERNARY	Qel	Altiplano
	Qlx	Brush Deposits and Lake Shores
	Qgm	Postglacial Marine Deposits
CRETACEOUS	Kfn	Fort Nelson
	Kfj	Fort St. John
DEVONIAN	Dfr	Town River
	Dh	Highway L.A.
	Dgu	Gunderson, Upper
	Dgm	Gunderson, Middle
	Ds	Gunderson, Lower
	Dh	Highway L.A.
	Dh	Highway L.A.
	Dh	Highway L.A.
	Dh	Highway L.A.
	Dh	Highway L.A.
	Dh	Highway L.A.
	Dh	Highway L.A.





CITIES SERVICE OIL CO.

PRELIMINARY PHOTOLOGICAL MAP
CANADA, NORTHWEST TERRITORIES
Pt. Providence Project
Kakisa River (W/2)


City Service Acreage	1000	1000
Local Photo Anomaly	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000
Photography - RCAF 1949, 1950 & 1951	1000	1000

Scale 1:250,000





White Lake 1:250,000 Sheet 65L
Photography - NCEP 1990



EXPLANATION

QUATERNARY

- Qal
Alluvium
- Qls
Beach Deposits and Lake Silts
- Qgm
Pleistocene Moraine Deposits

CRETACEOUS

- Kfn
Fort Nelson
- Kfsj
Fort St. John
- Kfsj
Fort St. John

DEVONIAN

UPPER

Dtr

Trout River

Dk

Kokosa Ls.

Dgw

Grumbler, Upper

Dgm

Grumbler, Middle

Do

Alexandra Fm.

Dh

Hay River Sh. & Ls.

Dsp

Slave Point (?) Reef

Dhr

Horn River Fm., Sh. & Ls. (Presqu'ile)


Opp

Pine Point Fm.

MIDDLE

Dfs

Fort Simpson



Local Photo Analysis

City Service Avenue


CITIES SERVICE OIL CO.

PRELIMINARY PHOTOGEOLOGIC MAP
CANADA, NORTHWEST TERRITORIES
Pt. Providence Project
White Lake

[illegible]

Copyright © 2004







60°00' 119°00'
Kakisa Quad (E/2) 1:250,000 Sheet 85D
Photography - RCAF 1948, 1950 & 1951

Local Photo Anomaly

EXPLANATION

QUATERNARY	Qal	Alluvium
	Qls	Beach Deposits and Lake Silts
	Qgm	Pleistocene Moraine Deposits
CRETACEOUS	Kfn	Fort Nelson
	Kfsj	Fort St. John
DEVONIAN	Dtr	Trout River
	Dk	Kakisa Ls.
	Dgu	Grumbler, Upper
	Dgm	Grumbler, Middle
	Da	Alexandra Fm.
	Dh	Hoy River Sh. & Ls.
	Dsp	Slope Point (?) Reef
	Dhr	Horn River Fm., Sh. & Ls. (Presquite)
	Dpp	Pine Point Fm.
	Dfs	Fort Simpson




CITIES SERVICE OIL CO.

PRELIMINARY PHOTOGEOLOGIC MAP
CANADA, NORTHWEST TERRITORIES
Ft. Providence Project
Kakisa River (E/2)

By Component	SCALE	1:250,000
Geology	DATE	June 1951
Topographic	GEOLGIST	S.S. Thorne
Photographic	PLOTTING	S.S. Thorne
Field Notes	DRAFTING	S.S. Thorne
Checked	CHECKED	S.S. Thorne









EXPLANATION

The diagram illustrates the geological column with three main periods: Quaternary, Cretaceous, and Devonian. Each period contains several geological units, some of which are grouped together or have specific stratigraphic relationships indicated by brackets and lines.

- QUATERNARY**
 - Qol (Alluvium)
 - Qls (Beach Deposits and Lake Silts)
 - Qgm (Pleistocene Marine Deposits)
- CRETACEOUS**
 - Kfn (Fort Nelson)
 - Kfnj (Fort St. John)
- DEVONIAN**
 - UPPER**
 - Dtr (Trout River)
 - Dk (Kekico Ls.)
 - Dgu (Gumbler, Upper)
 - Dgm (Gumbler, Middle)
 - Da (Alexander Fm.)
 - Dh (Hay River Fm., Sh. & Ls.)
 - Dap (Sage River (S) Reef)
 - Dtr (Hors River Fm., Sh. & Ls. (Shenep/Ts))
 - Dgo (Fox, coal Rn.)
 - MIDDLE**
 - Dfs (Fort Simpson)

CITIES SERVICE OIL CO.


PRELIMINARY PHOTOGEOLOGIC MAP
CANADA, NORTHWEST TERRITORIES
Ft. Providence Project
Folize Lake Quad W/2

Slip Component	SCALE	1/2 INCH
Slip Strike	047100	Normal, <u>NEO</u>
Vegetation Type	047100	S.S. <u>Forest</u>
Vegetation boundary	047100	S.S. <u>Forest</u>
Shrub Rock (log)	047100	S.S. <u>Forest</u>
Flats	047100	S.S. <u>Forest</u>
Infused Flats or Shrub	047100	S.S. <u>Forest</u>
Shrub	047100	S.S. <u>Forest</u>
Vegetation boundary	047100	S.S. <u>Forest</u>
Flats	047100	S.S. <u>Forest</u>
Threat Flats	047100	S.S. <u>Forest</u>
Slip - Shrub	047100	S.S. <u>Forest</u>
Surface (log) (Slip - Shrub)	047100	S.S. <u>Forest</u>



Time (h)	Control (n)	Treated (n)
0	100	100
15	95	90
30	90	85
45	85	80
60	80	75
75	75	70
90	70	65
105	65	60
120	60	40





Tathlina Quad (NW/4) 1:250,000 Sheet 85C
Photography - RCAF 1948, 1950 & 1951


EXPLANATION

QUATERNARY	Qal	CRETACEOUS	Kfn	DEVONIAN	Dtr
	Alluvium		Fort Nelson		Trout River
	Qls		Kfsj		D2l
	Beach Deposits and Lake Silts		Fort St. John		Dgu
	Qgm			UPPER	Dgm
	Pleistocene Moraine Deposits				Grumbler, Upper
					Dg
					Grumbler, Middle
				MIDDLE	Dh
					Alexandra Fm.
					Dhr
					Horn River Fm., Sh. & Ls.
					Dpp
					Stave Point (?) Reef
					Dpp
					Pine Point Fm.

Local Photo Anomaly

CITIES SERVICE OIL CO.

PRELIMINARY PHOTOGEOLGIC MAP
CANADA, NORTHWEST TERRITORIES
Ft. Providence Project
Tathlina Lake NW/4



Dip Component
Dip Strike
Topographic Slope
Formation boundaries
Strike fault lines
Faults
Inferred faults or structures
Elevation less than
Vegetation boundary
Bluffs
Thrust Fault
Dike - Bedrock Outcrop
Surface Color - Lt Gy (Light Grey)

SCALE 1:250,000
DATE November, 1960
GEOLOGY E.E. Thibault
PLOTTER E.E. Thibault
DRAFTING S.G.R.
CHECKED E.E. Thibault

Graphic Scale
0 1 2 3 4 5 Miles



