


Strategic Oil & Gas Ltd.

Work Order-Ref #: 19134

Vapor Intrusion Assessment (VIA) Soils Outside Casing (AGM)

Strategic et al Cameron
M-73

September 10 & 11, 2019



FORENSIC SOLUTIONS FOR ENERGY CHALLENGES

GCHEM Ltd.
BAY #1
4810-62ND AVE.
LLOYDMINSTER, AB
T9V 2E9
(780) 871-4668
www.gchem.ca
info@gchem.ca

Table of Contents

1.0	Executive Summary	4
2.0	Vapor Intrusions Assessment (VIA) Summary	5
2.1	Production Casing Assessment Summary	5
2.2	Surface Casing Vent Flow (SCVF) Assessment Summary	5
2.3	Soils Outside Casing (AGM) Assessment Summary	6
2.4	Interpreted Source of Migrating Gases	7
3.0	Background of Vapor Intrusion Assessments (VIA) at Resource Wells and Tracing Gas contents in the Environment using Energy Forensics	8
3.1	Biogenically Derived Methane Gas	9
3.2	Thermogenic Methane Gas	9
3.3	Classification, Characterization, and Geologic Origins of Combustible Gases in the Environment	9
4.0	Methods and Results	11
4.1	Field Assessment Methods	11
4.1.1	Non-Intrusive Vapor Intrusion Assessment	11
4.1.2	Intrusive Vapor Intrusion Assessment	13
4.2	Analytical Methods	15
5.0	Geochemical Measurements- Laboratory Results	16
5.1	Gases Obtained from Soil Vapor Probes (SVPs)	16
6.0	Conclusions	22

List of Tables

Table 1	Non-Intrusive Surface PMD	12
Table 2	Intrusive AGM-Soil Vapor Probes (SVPs)	14
Table 3	Gas components, isotopic compositions measured and the analytical error of measurements at GCHEM'S Analytical Laboratory	15
Table 4	High resolution molecular and stable carbon isotopic compositions of gases collected during this investigation	17

List of Figures

Figure 1	Non-Intrusive Surface Methane Scan (PMD)	12
Figure 2	Intrusive Methane Readings from Soil Vapor Probes	14
Figure 3	ΣC_{2+} vs Methane	18
Figure 4	ΣC_{6+} vs Ethane	19
Figure 5	$CH_4/\Sigma C_{2+}$ vs $\delta^{13}C$ CH_4	20
Figure 6	$\delta^{13}C$ CO_2 vs $\delta^{13}C$ CH_4	21

List of Attachments

Attachment 1	Well Site Photographs	23
Attachment 2	Chain of Custody	27
Attachment 3	Gas Sample Container Photographs	29
Attachment 4	Gas Analysis Data Sheets (GADS)	33

1.0 Executive Summary

On September 10-11, 2019, Strategic et al Cameron Hills M-73 was investigated for natural gas leakage in soils outside of casing. Total combustible gas (%LEL) and H₂S field instruments were used to measure gas levels and types at soil test sites. Gas samples were collected, contained and preserved from soils outside casing (AGM) and for baseline comparison, background locations, ~30m away from the wellbore were also assessed.

Soils outside casing immediately adjacent to and S-SW of the well bore were water saturated however, no gas bubbling was observed in standing water. A non-intrusive surface CH₄ scan was conducted in soils outside casing and at 3-background locations (N35m, W35m, and SW30m). Five sites (E3-S5, E2-S5, E2-S4, E2-S1, and S2-W1) contained elevated (22, 102, 15, 16 and 11 ppm v/v, respectively) above background methane readings (5 ppm v/v). All other sites tested near the wellbore contained CH₄ levels ranging from 2 to 4 ppm v/v and were similar to the BKG sites. A total of 11 soil sites outside casing were assessed for gas leakage using an intrusive methodology where 5cm vapor test holes were augered into soils and Soil Vapor Probes (SVPs) were inserted into each test hole. Of the 11 soil vapor test sites outside casing, 7 sites (N2m 238 ppm v/v, N4m 267 ppm v/v, E2m 126 ppm v/v, SE5m 625 ppm v/v, S1m 795 ppm v/v, W1m 1385 ppm v/v, and W3m 424 ppm v/v) contained elevated, above background (BKG N35m 35 ppm v/v) methane contents. SOG selected 2 soil sites (SE5m and W1m) to measure chemical and δ¹³C isotopic compositions to aid in classifying combustible gas contents. The 2-sites contained high levels of CH₄ gas and low, similar to background levels of associated ΣC₂₊ gas. Methane index values suggest that elevated %LEL values measured at the 2-soil sites suggest biogenic in origins for CH₄ gas. δ¹³C CH₄ and CO₂ values for gases obtained from test site W1m are consistent with methane generated via bacterial fermentation processes while the gases obtained from test site SE5m have been significantly altered by bacterial oxidation processes. However, low associated ΣC₂₊ gas levels support bacterial fermentation origins for CH₄ gas. ΣC₂₊ gas levels in soils outside casing at test sites SE5m and W1m are low, are similar to background and the result of natural movement of light hydrocarbon gases from reservoirs at depth, upward through subsurface fractures and micro-fractures to surface. This is a naturally occurring process in all hydrocarbon sedimentary basins in the world.

With information available to date, soil vapor test sites SE5m and W1m would be classified as 'Biogenic-Naturally Occurring CH₄-Non-Impacted'.

2.0 Vapor Intrusion Assessment Summary

Operating Company: Strategic Oil & Gas Ltd.
Well Name: Strategic et al Cameron M-73
UWI: 300M736010117150

License Number: 001765
Test Date September 10-11, 2019
GCHEM Project Number 19134

2.1 Production Casing Test Summary

Combustible Gas (CH₄) ([%LEL])	nm		
Hydrogen Sulphide (H₂S) Gas (ppm v/v)	nm		
PC Flow Rate (m³/day)	nm		
P-T Date Logger Installed	nm		
P-T Data Logger Removed	nm		
P-T Data Logger Test Duration	nm		
MAX Pressure (kPa)	nm		
Gas Spls. Collection-Measurement	Total Collected	Analysis Requested*	Classification**
PC Samples (Total)	0		
PC Combustible Gas Class. Level-1 (Chemical)		NA	NA
PC Combustible Gas Class. Level-2 ($\delta^{13}\text{C}$)		NA	NA
PC Combustible Gas Class. Level-3 (δD)		NA	NA
PC Combustible Gas Class. Level-4 (C^{14})		NA	NA

2.2 Surface Casing Vent Flow (SCVF) Test Summary

SCV Ten-Minute Bubble Test Result	nm		
SCV Flow Rate (m³/day)	nm		
SCVF Pressure-Temp Logger Installed	nm		
SCV Pressure-Temp Data Logger Removed	nm		
SCV Shut-In Time (hrs)	nm		
SCV MAX-Recorded Build Up Pressure (kPa)	nm		
SCV Stabilized Build-up Pressure (kPa):	nm		
SCV Stabilized Build-up Time (hours)	nm		
SCV Standpipe Max CH₄ Content (% LEL):	nm		
SCV Standpipe Max H₂S Content	nm		
SCV Gas Spls. Collection-Measurement	Total Collected	Analysis Requested*	Classification**
SCV Samples (Total)	0		
SCV Combustible Gas Class. Level-1 (Chemical)		NA	NA
SCV Combustible Gas Class. Level-2 ($\delta^{13}\text{C}$)		NA	NA
SCV Combustible Gas Class. Level-3 (δD)		NA	NA
SCV Combustible Gas Class. Level-4 (C^{14})		NA	NA

2.3 Soil Gas Migration Test – Soils Outside Casing (AGM) Summary

A) Non-Intrusive CH₄ Surface Soil Scan (PMD) (Figure 1 and Table 1)

Well Casing Surface CH ₄ Test Sites	105
MAX Surface CH ₄ Reading	102
MAX H ₂ S Well Soil Reading (ppm v/v)	<1
Number of Background Sites	3
MAX Background CH ₄ (ppm v/v)	5
Max H ₂ S BKG Soil Reading (ppm v/v)	<1
Surface CH ₄ -PMD Gas Classification	

B) Non-Intrusive Surface Enclosed Soil Vapor FLUX Chamber Test

Surface SV-FC CH ₄ Test Sites	1		
MAX SV-FC CH ₄ Reading	102		
SV-FC Gas Spls. Collection-Measurement	Total Collected	Analysis Requested*	Test Site
SV-FC Samples (Total)	2	Soil Vapor Flux not requested by client	
SV-FC & Sites Requested for Level-1 Analysis		NA	NA
Combustible Gas Classification Level-1 (Chem.)			NA
SV-FC & Sites Requested for Level-2 Analysis		NA	NA
Combustible Gas Classification Level-2 ($\delta^{13}\text{C}$)			NA
SV-FC & Sites Requested for Level-3 Analysis		NA	NA
Combustible Gas Classification Level-3 (δD)			NA
SV-FC & Sites Requested for Level-4 Analysis		NA	NA
Combustible Gas Classification Level-4 (^{14}C)			NA

C) Intrusive Auger Test Holes with Soil Vapor Probes (Figure 2 and Table 2).

Number Soil Vapor Probe (SVP) Test Sites	11		
MAX SVP CH ₄ Reading (%LEL)	2.8		
Max H ₂ S SVP Field Reading (ppm v/v)	<1		
Number SVP BKG Test Sites	3		
MAX SVP CH ₄ BKG Test Sites (ppm v/v)	35		
SVPs Gas Spl. Collection & Measurement	Total Collected	Analysis Requested*	Test Site
Soil Vapor Probes (SVPs) AGM (Total)	5		
SVPs & Sites Requested for Level-1 Analysis		2	SE5m, W1m
Combustible Gas Classification Level-1 (Chem.)			Biogenic, Non-Impacted
SVPs & Sites Requested for Level-2 Analysis		2	SE5m, W1m
Combustible Gas Classification Level-2 ($\delta^{13}\text{C}$)			Biogenic, Non-Impacted
SVPs & Sites Requested for Level-3 Analysis		0	NA
Combustible Gas Classification Level-3 (δD)			NA
SVPs & Sites Requested for Level-4 Analysis		0	NA
Combustible Gas Classification Level-4 (^{14}C)			NA



BKG Gas Spl. Collection-Measurement	Total Collected	Analysis Requested*	Test Site
BKG Soil Vapor Probe (SVPs) (Total)	3		
SVPs & Sites Requested for Level-1 Analysis		1	BKG N35m
Combustible Gas Classification Level-1 (Chem.)			Biogenic-Natural Occurring-Baseline
SVPs & Sites Requested for Level-2 Analysis		0	
Combustible Gas Classification Level-2 ($\delta^{13}\text{C}$)			NA
SVPs & Sites Requested for Level-3 Analysis		0	
Combustible Gas Classification Level-3 (δD)			NA
SVPs & Sites Requested for Level-4 Analysis		0	
Combustible Gas Classification Level-4 ($\text{^{14}\text{C}}$)			NA

* Sample selection for chemical and isotopic analysis (geochemical analytical suite) selected by client/operator.

2.4 Interpreted Source of Migrating Gases (measured depth from KB of the well)

Sample Point	Geologic Formation	Depth Range	Source Depth
SVP SE5m & W1m	Near Surface Soil Respiration		Biogenic CH ₄ , Non-Impacted, Baseline



3.0 Background of Vapor Intrusion Assessments (VIA) at Resource Wells & Tracing Gas Contents in the Environment using Energy Forensics

Undesired natural gas leakage from depth to surface at resource wells is becoming increasingly recognized and is a significant financial burden to the resource industry. When high levels of natural gas are found in the surface casing vent it is termed surface casing vent flow (SCVF) and when found in soils outside casing it is termed active gas migration (AGM). Identifying the source of leaking gas, maintaining zonal isolation and eliminating gas leakage to surface has proven to be a challenging task. Industry success rates using conventional gas leakage identification tools (e.g. noise, temperature, cement bond-integrity, ultra-sonic imaging logs, etc.) to eliminate surface gas migration in the first attempt is approximately 15% to 20%. Since 1997, through collaboration with industry, government regulators and academic institutions, GCHEM Ltd. has developed ‘Energy-Forensics’ and has obtained extensive expertise in field testing, gas sampling and preservation, analytical and interpretational techniques to pinpoint the geologic source of natural gases at resource wells.


It is important to note that detection of elevated combustible gases at surface does not always mean the well is impacted with deep sourced natural gas (thermogenic). Accurate gas characterization at well sites is critical as elevated CH₄ (%LEL) contents measured at or near surface may not indicate it is leaking or impacted (false-positive) but rather the combustible gases present are the result of biogenic activity or hydrocarbon contamination (or a combination of).

Thermogenic hydrocarbon gases have unique chemical and isotopic signatures based on many variables including the starting organic material they are produced from, the chemical processes from organic origin to current form, interaction with surrounding formation rock and fluids, and effects from migrating from origin to current trap. For example, molecular and isotopic composition ($\delta^{13}\text{C}$ and $\delta^2\text{H}$) of a low temperature, shallow sourced natural gas is significantly different with respect to those of a high temperature deep sourced natural gas. This principle allows the geologic source of leaking natural gas at a wellbore to be determined.


3.1 Biogenically Derived Methane Gas

As a normal part of soil respiration, methane may be generated via two biotic pathways (depending on substrate availability):

CO₂ reduction reaction



Fermentation Process



Biogenic methane gas may be further oxidized by bacteria. Oxidizing bacteria in soils preferentially consume ¹²C over ¹³C resulting the residual gas enriched in ¹³C (i.e. $\delta^{13}\text{C}$ values become less negative) with respect to the biogenic gas (isotope enriching effect). Therefore, biogenic oxidation may provide the false signature of a ‘mixture of biogenic and thermogenic methane’ or ‘thermogenic’ gas (GCHEM in prep). Biogenic CH₄ is generally prevalent in landfill or swamp gas.

3.2 Thermogenic Methane Gas

Methane gas can be generated by abiotic processes such as the thermo-degradation of organic matter at high pressure and temperature (thermogenesis). During thermogenic CH₄ generation, pending organic matter content, pressure and temperature, associated C₂+ gases may also be formed. Thermogenic CH₄ and C₂+ gases contain enriched $\delta^{13}\text{C}$ and δD values pending gas maturity, mixing and alteration and torturous pathway from source to trap.

3.3 Classification, Characterization and Geological Origins of Combustible Gases in the Environment.

Combustible gases in soils outside casing maybe classified and characterized (biogenic, thermogenic or mixed) using chemical, carbon and hydrogen and isotopic measurements and ¹⁴C concentrations. Leaking thermogenic natural gas in soils outside casing is easier to scientifically prove than biogenic methane sources. Elevated %LEL measured in AGM (on location) maybe the result of naturally occurring biogenic processes, anthropogenic leaking thermogenic natural gases and mixtures of both. A systematic 4-level approach can be used to determine the origins (biogenic-thermogenic or mixed) combustible gas contents and include:

1) Level-1 Characterization**High Resolution Compositional-Chemical Measurements.**

Permanent, inert and CH₄ to n-C₅H₁₂ & C₆+

See NGGC-1 CH₄ vs ΣC₂+ (Szatkowski et al 2000 & 2001).

See NGGC-2 C₂H₆ vs. C₆+ (Szatkowski et al 2000 & 2001).

2) Level-2 Characterization**Stable Carbon Isotope Measurements (δ¹³C).**

δ¹³C CH₄ to n-C₅H₁₂ & CO₂ (pending concentrations-gas levels).

See NGGC-3 CH₄/ΣC₂+ vs. δ¹³C CH₄ (Bernard 1978).

See NGGC-4 δ¹³C CO₂ vs. δ¹³C CH₄ (Whiticar 1993).

3) Level-3 Characterization**Hydrogen in Methane (δD).**

δD CH₄ to dD C₄H₁₂ (pending concentrations-gas levels).

See NGGC-5 δ¹³C CH₄ vs δD CH₄ (Coleman 1993).

4) Level-4 Characterization**¹⁴C pMC concentrations (radioactive ½ life of 5750 yr).**

Pending concentrations-gas levels.

¹⁴C reveals the age of the organic matter source from which CH₄ was generated but not the time of methanogenesis.

To determine the geological origins of leaking thermogenic natural gas contents, a series of plots developed by GCHEM Ltd are used and include.

1) Chemical & Isotopic Gas Field Diagram	C ₂ H ₆ /ΣC ₃ + vs δ ¹³ C C ₂ H ₆ (Szatkowski et al 2000, 2001).
2) Isotopic Gas Field Diagram	δ ¹³ C C ₂ H ₆ vs. δ ¹³ C C ₃ H ₈ (Szatkowski et al 2000, 2001).
3) Modified Chung Plot	δ ¹³ C vs 1/n (carbon & hydrogen number) (Chung 1988, and GCHEM Ltd. Unpublished).

Additional chemical and stable carbon and hydrogen isotopic plots have been developed to aid in determining the geological origins of natural gas found in the environment however, GCHEM has not published these novel and new correlations and relationships and they will not be shown or discussed in detail at this time (GCHEM Unpublished Internal Research).

4.0 Methods and Results



4.1 Field Assessment Methods and Results

4.1.1 Non-Intrusive Vapor Intrusion Assessment

On September 10, 2019, GCHEM conducted a surface soil methane scan using a Sensit PMD (Figure 1). Surface soil CH₄ levels were measured at 105 locations on a grid pattern (1m x 1m) covering approximately an 10m x 10m square area around the marked wellbore.

To establish background surface CH₄ gas levels a distance away from the well bore, three locations (35m north, 35m west, and 30m southwest of the wellbore) were also assessed. To enhance results of the surface methane scan and reduce potential effects from industrial contamination, at each test site, an atmospheric CH₄ gas level was recorded, the PMD gas sampling wand was coupled to surface soils and the CH₄ level was recorded for that specific test site. Atmospheric CH₄ level was subtracted from the CH₄ level measured after ground coupling to derive a surface soil CH₄ level at that point of the grid.

Five surface PMD sites (E3-S5, E2-S5, E2-S4, E2-S1, and S2-W1) contained elevated (22, 102, 15, 16 and 11 ppm v/v, respectively) above background (5 ppm v/v) CH₄ levels. All other surface sites assessed near the wellbore contained CH₄ levels that ranged from 2 to 4 ppm v/v that were similar to the BKG sites.

Figure 1. AGM Non-Intrusive Surface PMD

Figure-1A. NON-Intrusive CH₄ Surface Scan Well Casing Detail VIEW

Table 1. AGM Non-Intrusive Surface PMD


WELL CASING (AGM) Non-Intrusive Surface PMD (CH ₄) Soil Scan		WELL CASING (AGM) Non-Intrusive Surface PMD (CH ₄) Soil Scan		WELL CASING (AGM) Non-Intrusive Surface PMD (CH ₄) Soil Scan		WELL CASING (AGM) Non-Intrusive Surface PMD (CH ₄) Soil Scan	
Test Site (m)	PMD CH ₄ (ppm v/v) (%) Vol	Test Site (m)	PMD CH ₄ (ppm v/v) (%) Vol	Test Site (m)	PMD CH ₄ (ppm v/v) (%) Vol	Test Site (m)	PMD CH ₄ (ppm v/v) (%) Vol
N.5		E.5	2	S.5		W.5	
N1	2	E1	2	S1	4	W1	3
N2	2	E2	2	S2	4	W2	2
N3	2	E3	2	S3		W3	2
N4	2	E4	2	S4		W4	2
N5	2	E5	2	S5		W5	2
N5-E1	2	E5-S1	2	S5-W1		W5-N1	2
N4-E1	2	E5-S2	2	S4-W1		W5-N2	2
N3-E1	2	E5-S3	3	S3-W1	2	W5-N3	2
N2-E1	2	E5-S4	3	S2-W1	11	W5-N4	2
N1-E1	2	E5-S5		S1-W1		W5-N5	2
N1-E2	2	E4-S5	2	S1-W2		W4-N5	2
N2-E2	2	E4-S4	2	S2-W2		W4-N4	2
N3-E2	2	E4-S3	2	S3-W2		W4-N3	2
N4-E2	2	E4-S2	2	S4-W2		W4-N2	2
N5-E2	2	E4-S1	2	S5-W2		W4-N1	2
N5-E3	2	E3-S1	3	S5-W3		W3-N1	2
N4-E3	2	E3-S2	3	S4-W3		W3-N2	2
N3-E3	2	E3-S3	3	S3-W3	2	W3-N3	2
N2-E3	2	E3-S4	3	S2-W3	2	W3-N4	2
N1-E3	2	E3-S5	22	S1-W3	2	W3-N5	2
N1-E4	2	E2-S5	102	S1-W4	2	W2-N5	2
N2-E4	2	E2-S4	15	S2-W4	2	W2-N4	2
N3-E4	2	E2-S3		S3-W4	2	W2-N3	2
N4-E4	2	E2-S2		S4-W4	2	W2-N2	2
N5-E4	2	E2-S1	16	S5-W4	2	W2-N1	2
N5-E5	2	E1-S1	4	S5-W5	2	W1-N1	2
N4-E5	2	E1-S2	5	S4-W5	2	W1-N2	2
N3-E5	2	E1-S3		S3-W5	4	W1-N3	2
N2-E5	2	E1-S4		S2-W5	2	W1-N4	2
N1-E5	2	E1-S5		S1-W5	2	W1-N5	2

BACKGROUND Non-Intrusive Surface PMD (CH₄) Soil Scan

Test Site (m)	PMD CH ₄ (ppm v/v) (%)	Test Site (m)	PMD CH ₄ (ppm v/v) (%)	Test Site (m)	PMD CH ₄ (ppm v/v) (%)	Test Site (m)	PMD CH ₄ (ppm v/v) (%)
N35	2			SW30	3	W35	5

4.1.2 Intrusive Vapor Intrusion Assessment

A total of 11 soil sites outside casing were assessed for gas leakage using an intrusive methodology where soil vapor test holes were augered into soils and Soil Vapor Probes (SVPs) were inserted into each test hole. The SVPs were allowed to stabilize for approximately 30-minutes prior to combustible gas content measurement with the PMD (Figure 2). Of the 11 soil vapor test sites outside casing, 7 sites (N2 238 ppm v/v, N4 267 ppm v/v, E2 126 ppm v/v, SE5 625 ppm v/v, S1 795 ppm v/v, W1 1385 ppm v/v, and W3m 424 ppm v/v) contained elevated, above background (N35 35 ppm v/v) methane contents.

Figure 2. AGM Intrusive SVPs


Seeded grass surrounding the well casing is of similar stand-growth to surrounding lease vegetation. No stressed dead spots or discoloration was observed.

Table 2. AGM Intrusive SVPs

Intrusive AGM - Hand Auger-Test Hole-Install Soil Vapor Probes (SVPs) ATM-Isolated

Test Site (m)	Soil Vapor Probes		Soil Parameters		Gas		Site Assessment
	IR-CH ₄ (ppm v/v)	H ₂ S (ppm v/v)	Type	Moist. (1-5)	HC-CONT (Y-N)	Sample (Y-N)	Comments
N.5							Surface Water
N2	238	<1.0		5	no		
N4	267	<1.0		5	no		Yes
NE1.5	2	<1.0		5	no		
E.5							Surface Water
E2	126	<1.0		5	no		
E3	34	<1.0		5	no		
SE1.5	11	<1.0		5	no		
SE5	625	[1.3]		5	no	Yes	Final SVP Watered Out
S1	795	[1.6]		5	no	Yes	
S3							Surface Water
S5							Surface Water
W.5							Surface Water
W1	1385	[2.8]		5	no	Yes	
W3	424	<1.0		5	no	Yes	Final SVP Watered Out
NW1.5	16	<1.0		5	no	No	
Test Site (m)	Soil Vapor Probes		Soil Parameters		Gas		Site Assessment
	IR-CH ₄ (ppm v/v)	H ₂ S (ppm v/v)	Type	Moist. (1-5)	HC-CONT (Y-N)	Sample (Y-N)	Comments
BKG N35	35	<1.0		5	No	Yes	
BKG W35	2	<1.0		5	No	Yes	
BKG SW30	2	<1.0		5	No	Yes	

4.2 Analytical Methods

- a. High Resolution Compositional Analysis (HRCA).
 - i. He, H₂, O₂, N₂, CO₂, CH₄ to n-C₅H₁₂ & C₆+
- b. Stable Carbon ($\delta^{13}\text{C}$) and Hydrogen (δD) Isotopic Analysis.
 - i. $\delta^{13}\text{C}$ CH₄ to n-C₅H₁₂ and CO₂, and δD CH₄ to n-C₅H₁₂

Compositional (molecular) analyses were conducted at GCHEM's Analytical Laboratory using Hewlett Packard 5890 and Agilent 7890 Gas Chromatographs (GCs) configured for low (ppb v/v to ppm v/v) to high (vol. %) level detection of light alkane/alkene gases and atmospheric gas components. Chemical analysis of gases measured, and analytical error are shown in Table-1.

Stable carbon ($\delta^{13}\text{C}$) isotope ratios of light hydrocarbon gases (LHG) and carbon dioxide and hydrogen isotope ratios (δD) of LHG were also measured at GCHEM's Analytical Laboratory on a Thermo-Scientific MAT-253 Gas Chromatograph-Combustion-Continuous Flow-Isotope Ratio Mass Spectrometer (GC-C-CF-IRMS). Carbon isotope ratios are reported in delta (δ) notation and per mil (‰, parts per thousand) with respect to VPDB (Vienna Pee Dee Belemnite). Hydrogen isotope ratios are reported in delta (δ) notation and per mil (‰) with respect to VSMOW (Vienna Standard Mean Ocean Water).

Gas Component (ppmv)	Molecular Formula	Analytical Error (%)	Stable Carbon Isotopic Composition ($\delta^{13}\text{C}$)	Analytical Error (‰ VPDB)	Hydrogen Isotopic Composition (δD)	Analytical Error (‰ VSMOW)
Hydrogen	H ₂	±7%	-	-	$\delta\text{D H}_2$	±10
Helium	He	±7%	-	-	-	-
Nitrogen	N ₂	±7%	-	-	-	-
Oxygen	O ₂	±7%	-	-	-	-
Carbon Dioxide	CO ₂	±7%	$\delta^{13}\text{C CO}_2$	±0.2	-	-
Hydrogen Sulphide	H ₂ S	±7%	-	-	-	-
Methyl Mercaptan	CH ₄ S	±7%	-	-	-	-
Ethyl Mercaptan	C ₂ H ₆ S	±7%	-	-	-	-
Thiophene	C ₄ H ₄ S	±7%	-	-	-	-
Dimethyl Disulfide	C ₂ H ₆ S ₂	±7%	-	-	-	-
Methane	CH ₄	±7%	$\delta^{13}\text{C CH}_4$	±0.1	$\delta\text{D CH}_4$	±10
Ethane	C ₂ H ₆	±7%	$\delta^{13}\text{C C}_2\text{H}_6$	±0.2	$\delta\text{D C}_2\text{H}_6$	±10
Ethene	C ₂ H ₄	±7%	$\delta^{13}\text{C C}_2\text{H}_4$	±0.2	$\delta\text{D C}_2\text{H}_4$	±10
Propane	C ₃ H ₈	±7%	$\delta^{13}\text{C C}_3\text{H}_8$	±0.2	$\delta\text{D C}_3\text{H}_8$	±10
Propene	C ₃ H ₆	±7%	$\delta^{13}\text{C C}_3\text{H}_6$	±0.2	$\delta\text{D C}_3\text{H}_6$	±10
iso-Butane	i-C ₄ H ₁₀	±7%	$\delta^{13}\text{C i-C}_4\text{H}_{10}$	±0.2	$\delta\text{D i-C}_4\text{H}_{10}$	±10
normal-Butane	n-C ₄ H ₁₀	±7%	$\delta^{13}\text{C n-C}_4\text{H}_{10}$	±0.2	$\delta\text{D n-C}_4\text{H}_{10}$	±10
iso-Pentane	i-C ₅ H ₁₂	±7%	$\delta^{13}\text{C i-C}_5\text{H}_{12}$	±0.2	$\delta\text{D i-C}_5\text{H}_{12}$	±10
normal-Pentane	n-C ₅ H ₁₂	±7%	$\delta^{13}\text{C n-C}_5\text{H}_{12}$	±0.2	$\delta\text{D n-C}_5\text{H}_{12}$	±10
Hexane and higher	C ₆ +	±7%	-	-	-	-

Table 3. Gas components, isotopic compositions measured and the analytical error of the measurements at GCHEM's Analytical Laboratory.

5.0 Geochemical Measurements-Laboratory Results.

As part of this VIA (SCV-AGM), a total of 5 gas samples were collected, contained and preserved from the following locations or sample points: SVPs-soils outside casing (N4m, SE5m, S1m, W1m, and W3m) and 3-BKG locations (N35m, W35m and SW30m).

At the request of the Strategic Oil and Gas, chemical and $\delta^{13}\text{C}$ isotopic compositions were measured for gases obtained from two SVP test sites (SE5m and W1m) that contained elevated, above background, levels of combustible gases and chemical compositions for one BKG SVP (N35m). High Resolution chemical and $\delta^{13}\text{C}$ isotopic compositions were measured at GCHEM's Forensic Lab and are provided in Table 4.

5.1 Gases Obtained from Soil Vapor Probes (SVPs).

Gases measured in two SVPs test sites in soils near the well bore (SE5m and W1m) contain above atmospheric levels of CO₂ (3300 and 1276 ppm v/v, respectively). Methane gas was elevated (434.4 and 113.4 ppm v/v, respectively) when compared to background level measured at BKG N35 (32.46 ppm v/v) (Table 4 and Figure 3). C₂₊ gas levels in SVPs SE5m and W1m were low (<0.01) and similar to background levels (<0.01 ppm v/v). High methane with low, associated C₂₊ thermogenic gases suggests a biogenic or biotic source via CO₂ reduction or fermentation reactions for methane gas (Figures 5 and 6). C₆₊ gas contents at SVPs sites SE5m and W1m were low (0.18 and 0.06 ppm v/v respectively) and suggest hydrocarbon contamination was not present at SVP test sites (Figure 4).

Sufficient levels of C₂₊ gases at SVP sites SE5m and W1m were too low to measure $\delta^{13}\text{C}$ isotopic compositions. Sufficient levels of CO₂ and CH₄ were available for $\delta^{13}\text{C}$ at both sites. $\delta^{13}\text{C}$ CH₄ and $\delta^{13}\text{C}$ CO₂ at SVP SE5m was -61.85 and -17.23 ‰ VPDB, respectively. $\delta^{13}\text{C}$ CH₄ and $\delta^{13}\text{C}$ CO₂ at SVP W1m was -32.67 and -14.59 ‰ VPDB respectively (Table 4). $\delta^{13}\text{C}$ CH₄ and CO₂ values for gases at W1m are consistent with methane generated via bacterial fermentation processes while the gases at SE5m have been significantly altered by bacterial oxidation processes however, low, similar to background levels of $\Sigma\text{C2+}$ supports a biotic source for CH₄ gas via bacterial fermentation pathways (Figures 5 and 6).

Table 4. High resolution molecular and stable carbon isotopic compositions of gas samples collected as part of the VIA at Strategic et al Cameron M-73. Hydrogen isotopic compositions were not measured at the request of SOG.

Gas Component	Sample Point Date Collected	W1 Sept. 10-19 (ppm v/v)	SE5 Sept. 10-19 (ppm v/v)	BKG N35 Sept. 11-19 (ppm v/v)
Hydrogen		3.96	4.08	5.53
Helium		2.82	2.77	2.85
Nitrogen		777147	777640	775047
Oxygen		221457	218618	220518
Carbon Dioxide		1276	3300	4394
Methane		113.4	434.4	32.46
Ethane		<0.01	<0.01	<0.01
Ethene		<0.01	<0.01	<0.01
Propane		<0.01	<0.01	<0.01
Propene		<0.01	<0.01	<0.01
iso-Butane		<0.01	<0.01	<0.01
n-Butane		<0.01	<0.01	<0.01
iso-Pentane		0.13	0.20	<0.01
n-Pentane		<0.01	<0.01	<0.01
C ₆ +		0.06	0.18	0.09
<hr/>				
C1 Index (C1/ΣC2+)		11340	434400	3246
C2 Index (C2/ΣC3+)		N/A	N/A	N/A
C3 Index (C3/ΣC4+)		N/A	N/A	N/A
C4 Index (C4/C5)		N/A	N/A	N/A
ΣC2+		N/A	N/A	N/A
ATM Ratio (N2/O2)		3.51	3.56	3.51
Vol % CO ₂ of TG		0.13	0.33	0.44
Vol % Lt. Alk. of TG		0.01	0.04	0.00
Vol % Lt. Alk. CH4		99.88	100.0	100.0
Vol % Lt. Alk. C2+		0.12	0.05	0.00
Vol % C2+ of TG		0.00	0.00	0.00
<hr/>				
Stable Carbon Isotope Compositions (‰ VPDB)				
δ ¹³ C CH ₄		-32.67	-61.85	nm
δ ¹³ C C ₂ H ₆		nm	nm	nm
δ ¹³ C C ₂ H ₄		nm	nm	nm
δ ¹³ C C ₃ H ₈		nm	nm	nm
δ ¹³ C C ₃ H ₆		nm	nm	nm
δ ¹³ C i-C ₄ H ₁₀		nm	nm	nm
δ ¹³ C n-C ₄ H ₁₀		nm	nm	nm
δ ¹³ C i-C ₅ H ₁₂		nm	nm	nm
δ ¹³ C n-C ₅ H ₁₂		nm	nm	nm
δ ¹³ C CO ₂		-14.59	-17.23	nm
<hr/>				
Stable Hydrogen Isotopic Compositions (‰ VSMOW)				
δD H ₂		nm	nm	nm
δD CH ₄		nm	nm	nm
δD C ₂ H ₆		nm	nm	nm
δD C ₃ H ₈		nm	nm	nm
δD i-C ₄ H ₁₀		nm	nm	nm
δD n-C ₄ H ₁₀		nm	nm	nm
<hr/>				
14C Concentration (pMC)		nm	nm	nm




Figure-3. ΣC_2^+ vs Methane. Combustible gases detected in soils and SCVs at a wellhead may result from several origins. Natural gases indicative of SCVF or AGM are thermogenic in origin (natural gas in deep reservoirs), contain high methane and C_2^+ contents and plot in the Upper RH Quadrant. Low natural gas levels in background, off lease areas are naturally present in soils, vary from region to region and plot in the Lower LH Quadrant. Biogenic gases (swamp-gas) are produced by bacteria, are comprised of predominantly methane and plot in Lower RH Quadrant. Samples plotting in the Lower LH and RH do not contain SCVF or AGM and would not require down-hole remediation.




Figure 4. ΣC_6^+ vs Ethane. C_6^+ gases are relatively large molecules that do not readily or easily migrate in large quantities from depth upwards through subsurface fractures or micro-fractures to surface. Contamination by oil spills, fuels, and solvents is indicated by soil vapor samples that have high contents of C_6^+ compounds and plot in the Lower RH Quadrant. Samples plotting in the Lower LH and RH Quadrants do not contain evidence of either SCVF or AGM and would not require downhole repair operations.




Figure 5. $\text{CH}_4 / \Sigma\text{C}_2^+$ vs. $\delta^{13}\text{C CH}_4$. Thermogenic methane or methane generated by abiotic processes such as the thermal degradation of organic matter at high temperature and pressure (thermogenesis) contains enriched (less negative) $\delta^{13}\text{C}$ values ranging from -50 to -20‰ VPDB and methane relative to C_2^+ gas contents (gas wetness) less than 100. Methane gas can be generated by biotic processes such as the degradation of organic matter via CO_2 reduction or fermentation reactions generating biogenic methane. It should be noted that as a normal part of soil respiration, methane may be generated or destroyed by variable biotic pathways. Biogenic methane gas may be oxidized by bacteria resulting in an ‘isotopic enriching effect’ (i.e. $\delta^{13}\text{C}$ values become less negative as a result of oxidizing bacteria in soils that preferentially consume ^{12}C over ^{13}C , leaving the remaining gas enriched in ^{13}C). Since biogenic oxidation decreases the ratio between ^{12}C and ^{13}C , it may result in enriched $\delta^{13}\text{C CH}_4$ values that overlap with the MIXING or THERMOGENIC-GAS TREND. Biogenic methane may therefore contain $\delta^{13}\text{C}$ values greater than -50‰ VPDB (GCHEM Internal RD).




Figure 6 $\delta^{13}\text{C CO}_2$ vs. $\delta^{13}\text{C CH}_4$. Thermogenic methane or methane generated by abiotic processes such as the degradation of organic matter at high temperature and pressure contains enriched (less negative) $\delta^{13}\text{C}$ values ranging from -55 to -20‰ VPDB (or higher) and $\delta^{13}\text{C CO}_2$ values in the range of -25 to 4‰ VPDB. Methane gas may be generated by biotic processes such as the degradation of organic matter via CO_2 reduction or fermentation reactions generating biogenic methane. Biogenic methane may contain $\delta^{13}\text{C}$ values greater than -40‰ VPDB due to biogenic oxidation processes (GCHEM, in prep).

6.0 Conclusions

Soils outside casing are wet increasing the difficulty leaking natural gas assessments. 7 of the 11 intrusive soil sites outside casing tested for combustible gas contents contained elevated methane levels that ranged from 126 to 1385 ppm v/v. H₂S was not detected (< 1.0 ppm v/v) at any of the soil test hole sites. SOG selected SVP test sites SE5m (625 ppm v/v) and W1m (1385 ppm v/v) for high resolution chemical and stable carbon isotope measurements to classify combustible gas contents. Light hydrocarbon gases were dominated by methane gas while associated C₂+ gases were low and similar to background levels measured at test site BKG N35m. Methane index ratio (gas wetness) suggest that elevated %LEL values and associated CH₄ gas measured at the 2-soil sites are biogenic in origin. $\delta^{13}\text{C}$ CH₄ and $\delta^{13}\text{C}$ CO₂ isotopic compositions for gases obtained from SVP test sites W1m are consistent with methane generated via bacterial fermentation processes while the gases measured at E5m are enriched but have undergone primary bacterial oxidation and secondary alteration processes.

With information available to date, SVP soil test sites SE5m and W1m would be classified as ‘biogenic-baseline’ where CH₄ gas is the result of natural soil respirations processes via CO₂ reduction or fermentation processes generating biogenic CH₄. C₂+ gases in soils near the well are low, similar to background levels and the result of natural movement of thermogenic natural gas, from reservoirs at depth, upward through fractures and micro-fractures to surface. This is a naturally occurring process prevalent in every hydrocarbon sedimentary basin in the world.

Attachment-1

Strategic Oil & Gas Ltd.

Strategic Cameron M-73

Well Site Photographs







Attachment-2

Strategic Oil & Gas Ltd.

Strategic Cameron M-73

Chain of Custody (COC)



CHAIN OF CUSTODY

Box #1, 4810 62nd Ave., Daybreakmaster, Alberta T5V 2C9
E: info@gehrm.ca
Tel: (780) 871-4648
Fax: (780) 809-2885

GCHEM LTD.

GCHEM Ltd. Project# 78D

Client Information

Company Strategic Address _____
City, Prov. _____ Postal Code _____
Client Contact _____
Phone # _____ Fax # _____
E-Mail _____

Billing/Report Information

Analysis To Be Performed

Services Needed (TAT)

- Standard 5-7 Days ✓
- Rush 48hrs. _____
- priority Rush 24hrs. _____

(*) Working Days
(**) Call for Pricing and Advance Notice

1000

卷之三

Comments

Relinquished By: Will Rose Date/Time: Sept 16/19 Relinquished To: _____
Relinquished By: _____ Date/Time: _____ Relinquished To: _____

Date/Time: _____



Attachment-3

Strategic Oil & Gas Ltd.

Strategic Cameron M-73

Gas Sample Containers


Photographs

(red boxed pictures are samples measured for chemical and/or isotopic compositions)





N4m
September 10, 2019



S1m
September 10, 2019




W1m
September 10, 2019



W3m
September 10, 2019



SE5m
September 10, 2019



E2-S5 Chamber
September 10, 2019



N4m
September 11, 2019



W1m
September 11, 2019



SE5m
September 11, 2019



SE5m Flux
September 11, 2019



BKG N35m
September 11, 2019





Attachment-4

Strategic Oil & Gas Ltd.

Strategic Cameron M-73

Gas Analysis Data Sheets (GADS)

 GCHEM LTD.		HIGH RESOLUTION GAS ANALYSIS CARBON ISOTOPE ANALYSIS HYDROGEN ISOTOPE ANALYSIS				
		Sampling Company	GCHEM Ltd.	Lab Sample No.	19134-06	
Date Tested	September 10, 2019	Test Type	Soil gas			
Operator Name	Strategic Oil & Gas	Sample Container Type	Glass Bottle			
Unique Well Identifier	M-73	Sampling Point	W1			
Well Name	not provided	Test Intervals or Perfs mKB	N/A			
Field or Area	not provided	Date Received	September 16, 2019			
Pool or Zone	not provided	Date Reported	October 3, 2019			
Well License	not provided	Entered By	Xiaolong Wang			
H2S Level (Observed at Site)	not provided	Reviewed By	Brad Johnston			
Sample Handling Conditions						
	Source/Sampled	Received				
Pressure (kPa)	N/A	60				
Temperature (°C)	N/A	20				
Other Information:						
Laboratory Analysis						
Component	HRGC Analysis As Received Mol Frac.	Air Free As received Mol Frac.	Air Free / Acid Free As Received Mol Frac.	Carbon Isotope Analysis ‰ VPDB	Hydrogen Isotope Analysis ‰ VSMOW	HRGC Analysis As Received ppm v/v
Neon	0.000014	0.009959	0.009959			13.90
Hydrogen	0.000004	0.002837	0.002837			3.96
Helium	0.000003	0.002021	0.002021			2.82
Nitrogen	0.777147	0.000000	0.000000			777147
Oxygen	0.221457	0.000010	0.000010			221457
Carbon Dioxide	0.001276	0.913784	0.913784	-14.59		1276
Carbonyl Sulphide	nm	nm	nm			nm
Hydrogen Sulphide	nm	nm	nm			nm
Methyl Mercaptan	nm	nm	nm			nm
Ethyl Mercaptan	nm	nm	nm			nm
Thiophene	nm	nm	nm			nm
Dimethyl Disulphide	nm	nm	nm			nm
Methane	0.000113	0.081208	0.081208	-32.67		113.4
Ethane	0.000000	0.000000	0.000000			<0.01
Ethene	0.000000	0.000000	0.000000			<0.01
Propane	0.000000	0.000000	0.000000			<0.01
Propene	0.000000	0.000000	0.000000			<0.01
iso-Butane	0.000000	0.000000	0.000000			<0.01
n-Butane	0.000000	0.000000	0.000000			<0.01
iso-Pentane	0.000000	0.000096	0.000096			0.13
n-Pentane	0.000000	0.000000	0.000000			<0.01
C ₆ +	0.000000	0.000045	0.000045			0.06
TOTAL	1.000000	1.000000	1.000000			1000000
Properties						
Compositional Indicies		Real Gross Heating Value (mj/m3) @15°C and 101.35 kPa		Relative Density		
Vol % Hydrocarbons	0.01	Air Free as received	Moisture and Acid Gas Free	Calc. Mol. Mass Ratio	Calc. Relative Density	
Vol % CH ₄	99.88	0.00	3.14	0.9984	0.9984	
Vol % C ₂ +	0.00	Pseudo Critical Properties				
CH ₄ / \sum C ₂ +	N/A	pPc (kPa)	As Received	Acid Gas Free		
C ₂ / \sum C ₃ +	N/A	3761	7124			
C ₃ / \sum n-C ₄	N/A	pTc (°K)	133	294		
Geological Origin of Natural Gas						
Geological Formation		Depth Range (MD from KB of Well)		Probable Depth (MD from KB of Well)		
Comments						
Forensic Solutions for Oilfield Challenges						
GCHEM Ltd. Bay #1, 4810-62 Avenue Lloydminster, AB T9V 2E9 Tel: (780) 871-4668 Fax: (780) 808-8883 e-mail: info@gchem.ca GPA 2145-09. Revision 1.3, August 1, 2016						



GCHEM LTD.

HIGH RESOLUTION GAS ANALYSIS CARBON ISOTOPE ANALYSIS HYDROGEN ISOTOPE ANALYSIS

Sampling Company	GCHEM Ltd.	Lab Sample No.	19134-03
Date Tested	September 10, 2019	Test Type	Soil gas
Operator Name	Strategic Oil & Gas	Sample Container Type	Glass Botte
Unique Well Identifier	M-73	Sampling Point	SE5
Well Name	not provided	Test Intervals or Perfs mKB	N/A
Field or Area	not provided	Date Received	September 16, 2019
Pool or Zone	not provided	Date Reported	October 3, 2019
Well License	not provided	Entered By	Xiaolong Wang
H2S Level (Observed at Site)	not provided	Reviewed By	Brad Johnston

Sample Handling Conditions

	Source/Sampled	Received
Pressure (kPa)	N/A	47
Temperature (°C)	N/A	20

Other Information:

Laboratory Analysis

Component	HRGC Analysis As Received Mol Frac.	Air Free As received Mol Frac.	Air Free / Acid Free As Received Mol Frac.	Carbon Isotope Analysis ‰ VPDB	Hydrogen Isotope Analysis ‰ VSMOW	HRGC Analysis As Received ppm v/v
Neon	0.000014	0.003723	0.003723			13.93
Hydrogen	0.000004	0.001091	0.001091			4.08
Helium	0.000003	0.000740	0.000740			2.77
Nitrogen	0.777640	0.000000	0.000000			777640
Oxygen	0.218618	0.000003	0.000003			218618
Carbon Dioxide	0.003300	0.881974	0.881974	-17.23		3300
Carbonyl Sulphide	nm	nm	nm			nm
Hydrogen Sulphide	nm	nm	nm			nm
Methyl Mercaptan	nm	nm	nm			nm
Ethyl Mercaptan	nm	nm	nm			nm
Thiophene	nm	nm	nm			nm
Dimethyl Disulphide	nm	nm	nm			nm
Methane	0.000434	0.116089	0.116089	-61.85		434.4
Ethane	0.000000	0.000000	0.000000			<0.01
Ethene	0.000000	0.000000	0.000000			<0.01
Propane	0.000000	0.000000	0.000000			<0.01
Propene	0.000000	0.000000	0.000000			<0.01
iso-Butane	0.000000	0.000000	0.000000			<0.01
n-Butane	0.000000	0.000000	0.000000			<0.01
iso-Pentane	0.000000	0.000055	0.000055			0.20
n-Pentane	0.000000	0.000000	0.000000			<0.01
C ₆ +	0.000000	0.000048	0.000048			0.18
TOTAL	1.000000	1.000000	1.000000			1000000

Properties

Compositional Indices		Real Gross Heating Value (mJ/m3) @15°C and 101.35 kPa		Relative Density	
Vol % Hydrocarbons	0.04	Air Free	Moisture and	Calc. Mol. Mass Ratio	Calc. Relative Density
Vol % CH ₄	99.95	as received	Acid Gas Free	0.9990	0.9990
Vol % C ₂ +	0.00				
CH ₄ / \sum C ₂ +	N/A	0.02	4.43		
C ₂ / \sum C ₃ +	N/A				
C ₃ / \sum n-C ₄ +	N/A				

Pseudo Critical Properties

	As Received	Acid Gas Free
pPc (kPa)	3765	7047
pTc (°K)	133	290


Geological Origin of Natural Gas

Geological Formation	Depth Range (MD from KB of Well)	Probable Depth (MD from KB of Well)

Comments

Forensic Solutions for Oilfield Challenges
GCHEM Ltd. Bay #1, 4810-62 Avenue Lloydminster, AB T9V 2E9 Tel: (780) 871-4668 Fax: (780) 808-8883 e-mail: info@gchem.ca www.gchem.ca GPA 2145-09. Revision 1.3, August 1, 2016



 GCHEM LTD.		HIGH RESOLUTION GAS ANALYSIS CARBON ISOTOPE ANALYSIS HYDROGEN ISOTOPE ANALYSIS			
		Sampling Company	GCHEM Ltd.	Lab Sample No.	19134-07
Date Tested	September 11, 2019	Test Type	Soil gas		
Operator Name	Strategic Oil & Gas	Sample Container Type	Glass Bottles		
Unique Well Identifier	M-73	Sampling Point	BKG N35		
Well Name	not provided	Test Intervals or Perfs mKB	N/A		
Field or Area	not provided	Date Received	September 16, 2019		
Pool or Zone	not provided	Date Reported	October 3, 2019		
Well License	not provided	Entered By	Xiaolong Wang		
H2S Level (Observed at Site)	not provided	Reviewed By	Brad Johnston		
Sample Handling Conditions					
	Source/Sampled	Received			
Pressure (kPa)	N/A	62			
Temperature (°C)	N/A	20			
Other Information:					
Laboratory Analysis					
Component	HRGC Analysis As Received Mol Frac.	Air Free As received Mol Frac.	Air Free / Acid Free As Received Mol Frac.	Carbon Isotope Analysis ‰ VPDB	Hydrogen Isotope Analysis ‰ VSMOW
Neon	0.000013	0.002970	0.002970		
Hydrogen	0.000006	0.001248	0.001248		
Helium	0.000003	0.000644	0.000644		
Nitrogen	0.775047	0.000000	0.000000		
Oxygen	0.220518	0.000003	0.000003		
Carbon Dioxide	0.004394	0.990766	0.990766		
Carbonyl Sulphide	nm	nm	nm		
Hydrogen Sulphide	nm	nm	nm		
Methyl Mercaptan	nm	nm	nm		
Ethyl Mercaptan	nm	nm	nm		
Thiophene	nm	nm	nm		
Dimethyl Disulphide	nm	nm	nm		
Methane	0.000032	0.007318	0.007318		
Ethane	0.000000	0.000000	0.000000		
Ethene	0.000000	0.000000	0.000000		
Propane	0.000000	0.000000	0.000000		
Propene	0.000000	0.000000	0.000000		
iso-Butane	0.000000	0.000000	0.000000		
n-Butane	0.000000	0.000000	0.000000		
iso-Pentane	0.000000	0.000000	0.000000		
n-Pentane	0.000000	0.000000	0.000000		
C ₆ +	0.000000	0.000021	0.000021		
TOTAL	1.000000	1.000000	1.000000		
1000000					
Properties					
Compositional Indicies		Real Gross Heating Value (mj/m ³) @15°C and 101.35 kPa		Relative Density	
Vol % Hydrocarbons	0.00	Air Free as received	Moisture and Acid Gas Free	Calc. Mol. Mass Ratio	Calc. Relative Density
Vol % CH ₄	100.00	0.00	0.30	1.0000	1.0000
Vol % C ₂ +	0.00	Pseudo Critical Properties			
CH ₄ / \sum C ₂ +	N/A	pPc (kPa)	As Received	Acid Gas Free	
C ₂ / \sum C ₃ +	N/A	3772		7349	
C ₃ / \sum n-C ₄ +	N/A	pTc (°K)	133	303	
Geological Origin of Natural Gas					
Geological Formation		Depth Range (MD from KB of Well)		Probable Depth (MD from KB of Well)	
Comments					
Forensic Solutions for Oilfield Challenges					
GCHEM Ltd. Bay #1, 4810-62 Avenue Lloydminster, AB T9V 2E9 Tel: (780) 871-4668 Fax: (780) 808-8883 e-mail: info@gchem.ca www.gchem.ca GPA 2145-09. Revision 1.3, August 1, 2016					